

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/112610/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Aubineau, Jérémie, El Albani, Abderrazak, Chi Fru, Ernest ORCID: <https://orcid.org/0000-0003-2673-0565>, Gingras, Murray, Batonneau, Yann, Buatois, Luis A., Geffroy, Claude, Labanowski, Jérôme, Laforest, Claude, Lemée, Laurent, Mángano, Maria G., Meunier, Alain, Pierson-Wickmann, Anne-Catherine, Recourt, Philippe, Riboulleau, Armelle, Trentesaux, Alain and Konhauser, Kurt O. 2018. Unusual microbial mat-related structural diversity 2.1 billion years ago and implications for the Francevillian biota. *Geobiology* 16 (5) , pp. 476-497. 10.1111/gbi.12296 file

Publishers page: <http://dx.doi.org/10.1111/gbi.12296>
<<http://dx.doi.org/10.1111/gbi.12296>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Unusual microbial mat-related structural diversity 2.1 billion**
2 **years ago and implications for the Francevillian biota**

3 Jérémie Aubineau¹, Abderrazak El Albani^{1*}, Ernest Chi Fru², Murray Gingras³, Yann
4 Batonneau¹, Luis A. Buatois⁴, Claude Geffroy¹, Jérôme Labanowski¹, Claude Laforest¹,
5 Laurent Lemée¹, M. Gabriela Mángano⁴, Alain Meunier¹, Anne-Catherine Pierson-
6 Wickmann⁵, Philippe Recourt⁶, Armelle Riboulleau⁶, Alain Trentesaux⁶ & Kurt Konhauser³

7
8 ¹UMR CNRS IC2MP 7285, University of Poitiers, 86073 Poitiers, France. ²Cardiff
9 University, Main Building, Room 2.15B Park Place Cardiff, CF10 3AT, UK. ³Department of
10 Earth and Atmospheric Sciences University of Alberta Edmonton, Alberta, T6G 2E3 Canada.
11 ⁴Department of Geological Sciences, University of Saskatchewan, Saskatoon, SK S7N 5A5,
12 Canada. ⁵Department Geosciences, UMR 6118, University of Rennes 1, 35042 Rennes,
13 France. ⁶UMR 8187 LOG CNRS, University of Lille, ULCO, 59655, Villeneuve d'Ascq,
14 France.

15
16 **Abstract**

17 The 2.1-billion-year-old (Ga) Francevillian series in Gabon hosts some of the oldest reported
18 macroscopic fossils of various sizes and shapes, stimulating new debates on the origin,
19 evolution and organization of early complex life. Here we document ten representative types
20 of exceptionally well-preserved mat-related structures, comprising ‘elephant-skin’ textures,
21 putative macro-tufted microbial mats, domal buildups, flat pyritized structures, discoidal
22 microbial colonies, horizontal mat growth patterns, wrinkle structures, ‘kinneyia’ structures,
23 linear patterns and nodule-like structures. A combination of petrographic analyses, scanning
24 electron microscopy, Raman spectroscopy and organic elemental analyses of carbon-rich
25 laminae and microtexture converge on a biological origin for these structures. The observed
26 microtextures encompass oriented grains, floating silt-sized quartz grains, concentrated heavy

minerals, randomly oriented clays, wavy-crinkly laminae and pyritized structures. Based on comparisons with modern analogues, as well as an average $\delta^{13}\text{C}$ organic matter (C_{org}) composition of $-32.94 \pm 1.17\text{‰}$ (1 standard deviation, s.d.) with an outlier of -41.26‰ , we argue that the mat-related structures contain relicts of multiple carbon pathways including heterotrophic recycling of photosynthetically derived C_{org} . Moreover, the relatively close association of the macroscopic fossil assemblages to the microbial mats may imply that microbial communities acted as potential benthic O_2 oases linked to oxyphototrophic cyanobacterial mats and grazing grounds. In addition, the mat's presence likely improved the preservation of the oldest large colonial organisms, as they are known to strongly biostabilize sediments. Our findings highlight the oldest community assemblage of microscopic and macroscopic biota in the aftermath of the 'Great Oxidation Event', widening our understanding of biological organization during Earth's middle age.

Introduction

The Paleoproterozoic Era hosted one of the most important geochemical events in Earth's history, marked by a measurable accumulation of atmospheric oxygen, the so-called 'Great Oxidation Event' (GOE) between 2.45–2.32 billion years ago (Ga). Evidence for the GOE comes from the disappearance of detrital pyrite, uraninite and siderite from fluvial and deltaic deposits, an increase in the retention of iron in paleosols, an enrichment of Cr and U in iron formations, and perhaps most importantly, the disappearance of sedimentary sulfur isotope mass-independent (S-MIF) anomalies indicative of atmospheric SO_2 processing in the absence of appreciable ozone (Holland, 2002; Bekker *et al.*, 2004; Farquhar *et al.*, 2011; Konhauser *et al.*, 2011; Partin *et al.*, 2013a; Lyons *et al.*, 2014). In the wake of the GOE, large positive excursions in the $\delta^{13}\text{C}$ of marine inorganic carbon, during the 'Lomagundi Event' *ca.* 2.22–2.1 Ga (Karhu & Holland, 1996), is believed to reflect large-scale burial of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

organic matter in marine sediments (Berner, 2004; Bekker & Holland, 2012). New evidence suggests that the ‘Lomagundi Event’ ended with a drastic drop in Earth’s oxygen content (*e.g.* Partin *et al.*, 2013b), perhaps related to large-scale oxidation of organic carbon (Canfield *et al.*, 2013) and/or changes in ocean biogeochemical processes that may have inhibited primary productivity (Chi Fru *et al.*, 2015).

The Francevillian sedimentary rocks *ca.* 2.1 Ga record these biogeochemical fluctuations in Earth’s oxygen accumulation dynamics through extreme excursions in the carbon cycle, in addition to hosting the oldest large colonial macroorganisms (El Albani *et al.*, 2010, 2014). Major elements, trace metals, organic carbon and isotope analyses offer constraints on the basin geochemistry, revealing variations from the base to the top that are linked to sea level changes (Canfield *et al.*, 2013; Bankole *et al.*, 2016). The Lower Francevillian fluvatile rocks were deposited in oxygenated waters (Bankole *et al.*, 2016), whereas the Upper Francevillian marine rocks were deposited in oxic, ferruginous and euxinic waters (Canfield *et al.*, 2013). The large macrofossils are highly variable in terms of size, shape and pyritization process being represented by lobate, elongated and rod-shaped as well as disk-shaped morphotypes. The pyritization process did not fully occur in the circular disks, indicating that their organic composition differs from that of pyritized specimens (El Albani *et al.*, 2010, 2014). In addition, some carbonaceous spheroidal microstructures have also been reported as organic-walled acritarchs that might have a planktonic origin.

Recently, microbial fossils were found in the Francevillian sequence, including putative bacteriomorphous structures (Dubois *et al.*, 2015) and a Gunflint-type assemblage of microfossils in the stromatolitic units (Lekele Baghekema *et al.*, 2017). The presence of colonial macroorganisms and microfossils makes the Francevillian biota unique, but interestingly, evidence of microbial mats has never been demonstrated. In this work, we describe for the first time ever microbial mat-related structures (MRS) from 2.1 Ga

1
2
3 77 siliciclastic sandstones and black shales from the FB2 Member of the Francevillian basin,
4
5 78 Gabon. MRS are physical remains of destroyed mats and structures associated with decay,
6
7 79 instead of well-preserved mat growth features (Eriksson *et al.*, 2010). Their sedimentary
8
9 80 features are preserved because microbes secrete extracellular polymeric substances (EPS), an
10
11 81 adhesive mucilage (Decho, 2000), that provide a coherent and protective coating that is
12
13 82 unlikely to arise in sediments without a biological input (Gerdes *et al.*, 1993, 2000; Porada &
14
15 83 Bouougri, 2007; Noffke, 2010).

16
17
18 84 Microbial life in the Paleoproterozoic has been poorly described (Davies *et al.*, 2016)
19
20 85 even though existing paleontological and biogeochemical evidence points towards the
21
22 86 existence of significant microbial diversity previously in the Archean. Through a combination
23
24 87 of macroscopic and microtexture analyses, petrographic, geochemical, organic elemental
25
26 88 analyses and stable carbon isotope composition of bulk organic matter, we compare modern
27
28 89 and fossil mat-related structures, in order to characterize the marine paleoenvironment in
29
30 90 which the putative microbial mats formed. The data reveal the biogenicity of these structures,
31
32 91 adding new insights into the origins of the Francevillian biota (El Albani *et al.*, 2010; Ossa
33
34 92 Ossa, 2010; Parize *et al.*, 2013; El Albani *et al.*, 2014; Ngombi Pemba, 2014; Reynaud *et al.*,
35
36 93 2017). Importantly, we assessed and described the organisational, ecological and
37
38 94 taphonomical processes that led to the development and preservation of the unique
39
40 95 Francevillian biota.
41
42
43
44 96

45 46 97 **Geological background**

47
48 98 The Francevillian basin, located in the southeastern part of the Republic of Gabon (Figure 1a),
49
50 99 is a 35,000 km² depression comprising 2.2-2.0 Ga marine sedimentary rocks that
51
52 100 unconformably overly an Archean basement (Weber, 1969). Petrographic and geochemical
53
54 101 analyses indicate that the Francevillian sediments were not affected by metamorphic
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

transformation (maximum temperature 100 °C; Gauthier-Lafaye & Weber, 1989; Ngombi-Pemba *et al.*, 2014), resulting in the preservation of seawater composition at the time when deposition took place. Moreover, carbon isotopic and redox sensitive element compositions suggest that the ‘Lomagundi Event’ was a global event (Canfield *et al.* 2013).

The basin fill, composed of siliciclastic fluvial and marine deposits that are 1000–2500 m thick, is subdivided into four major lithostratigraphic formations, FA to FD (Figure 1b; Weber, 1969; Gauthier-Lafaye & Weber, 1989; Gauthier-Lafaye, 2006; El Albani *et al.*, 2010, 2014). Fluvatile to deltaic conglomeratic sandstones, overlain by marine sandstones deposited in a tidal environment (i.e., syntectonic filling), make up the FA Formation, with the overlying sediments hosting diagenetic uranium ore deposits and the natural nuclear fission reactors of Oklo (Gauthier-Lafaye & Weber, 1989, 2003). The marine-dominated FB sequence rests unconformably on FA. The former was deposited below storm wave base during basin deepening, and is subdivided into the FB1 (a, b and c) and FB2 (a and b) subunits. The FB1a and FB1b subunits are composed of interbedded shales, sandstones and conglomerate, stacked into fining upwards packages, while the overlying FB1c subunit mainly consists of black shales with a thin iron formation, likely corresponding to a maximum flooding surface. This is then overlain by thick Mn-rich carbonates. Massive sandstone beds of the FB2a subunit, probably deposited in channels near the fair-weather wave base, are capped by thinly laminated black shales that are 5 m thick, and interbedded with thin siltstone layers (FB2b), presumably deposited by waning storm surges. Outcrops of the FB2b subunit are scarce but host the well-known colonial macroorganisms described in El Albani *et al.* (2010, 2014). The overlying FC Formation consists of dolomite and thickly-banded stromatolitic cherts that were deposited under shallow-water conditions (Bertrand-Sarfati & Potin, 1994). The uppermost FD Formation consists mainly of black shales, with pyroclastic material at the top deposited in a deep marine environment (Gauthier-Lafaye & Weber, 2003).

The Francevillian basin has been intensively studied because of economic interests in their uranium and manganese ore content (Gauthier-Lafaye & Weber, 1989, 2003). Consequently, various ages have been obtained for the Francevillian sediments. For example, monazites contained in Archean plutonic rocks from the Chaillu massif close to the FA Formation transition provided U-Pb ages of 2998 ± 25 Ma to 2621 ± 30 Ma (Mouélé *et al.*, 2014). A U-Pb age of 2050 ± 30 Ma reported for uranium mineralization, delineates early diagenesis at the FA-FB boundary (Gancarz, 1978). Coarse-grained syenites of the N’Goutou volcanic complex, in the northern part of the Republic of Gabon, that are interlayered with rocks at the base of the FB1 sequence, yield an Rb-Sr age of 2143 ± 143 Ma (Bonhomme *et al.*, 1982). It is considered that these volcanic rocks were formed simultaneously with the sedimentary rocks. Diagenetic illites from the top of the FB1b subunit have been dated at 2099 ± 115 Ma by using the Sm-Nd method (Bros *et al.*, 1992). Finally, recent dating of zircons from welded tuffs near the top of the FD Formation produced a U-Pb age of 2083 ± 6 Ma (Horie *et al.*, 2005).

The Paleoproterozoic Francevillian MRS were discovered in the Moulendé quarry, the same fossiliferous quarry where large colonial organisms have been reported (El Albani *et al.*, 2010, 2014). MRS are heterogeneously distributed in the Upper FB2a subunit, as well as in the overlying laminated black shale. The latter is 5 m thick and forms the FB2b subunit (Figure 2), covering a ~20 m thick host interval.

Methods

Sampling and sample preparation

About one hundred samples were collected in order to describe large-scale variability in morphology, structural organization and diversity from outcrops localized in one locality over several years (Figure 1). All samples were logged to provide information on their spatial and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

time repartitions of the studied area (Figure 2), and then photographed at the University of Poitiers using a Nikon Europe D610 digital single-lens reflex camera equipped with a Nikon AF-S 24-120 mm f/4G ED VR lens. Polished slabs were observed using a ZEISS Discovery.V8 stereoscope combined with Axio Cam ERc 5s microscope camera. Based on this pre-screening analysis, representative samples were selected and powdered in an agate mortar for elemental and carbon isotope systematics.

Petrographic and mineralogical analysis

Petrographic and mineralogical examinations were performed by transmitted and reflected light on eight polished thin sections using a Nikon ECLIPSE E600 POL microscope equipped with a Nikon Digital Sight DS-U1 camera and NIS-Element D software for scanning observations. Seven thin sections were carbon coated and examined for textural and compositional analyses using a FEI Quanta 200 scanning electron microscope (SEM) at the University of Lille 1. The SEM, coupled to Rontec energy-dispersive spectra (EDS) for semi-quantitative mineral analysis in backscattered electron mode (BSE), was operated at 15 kV accelerating volts and a 1 nA beam current at a working distance of 10.5 mm.

Carbon analysis

Raman spectroscopy was used to determine the composition of carbon preserved within the MRS on two representative samples. Analyses were carried out at the University of Poitiers with a HORIBA JOBIN YVON Labram HR800UV, an integrated Olympus confocal microscope coupled to a Peltier-cooled CCD detector. All analyses were performed by means of 514.5 nm Ar⁺ laser of 1 mW, 200 µm confocal hole, 1800 grooves/mm grating. Data recording and treatment were done with LabSpec 5 software.

Ten representative samples, composed of five mat samples and five associated

surrounding sediments, were crushed for the determination of the elemental content of associated organic matter, as well as carbon and sulfur composition, at the University of Poitiers. Analyses were performed using a CHNS analyser, model FlashEA 1112 (ThermoFisher Scientific) by flash dynamic combustion at 970 °C under a constant flow of helium. Data were recorded using Eager 300 software. A calibration curve was obtained with sulfanilamide and BBOT – 2,5-Bis(5-tert-butyl-benzoxazol-2-yl)thiophene – before each analysis. The results of carbon contents include both inorganic and organic carbon, but petrographic analyses showed that calcium carbonate and dolomite concentrations were low.

Stable carbon isotope measurements (^{12}C , ^{13}C) of organic matter from 14 mat samples (from both FB2a and FB2b subunits) were conducted at the Stable Isotopes Laboratory of UMR Pegase (INRA Rennes, France), using an elemental analyser coupled with an isotope ratio mass spectrometer (IRMS) (VG Isoprime; UMR PEGASE – INRA Rennes). More than 500 mg of fine-grained powders were analysed and loaded in tin capsules for each sample. IAEA-C6 (sucrose ANU, $\delta^{13}\text{C} = -10.63\text{‰}$), USGS-24 (graphite, $\delta^{13}\text{C} = -15.98\text{‰}$), USGS-40 (glutamic acid, $\delta^{13}\text{C} = -26.32\text{‰}$) and Urea Isotopic Working Standard ($\delta^{13}\text{C} = -38.3\text{‰}$) were used as reference standards. Data are expressed as delta values (‰) relative to Vienna Pee Dee Belemnite (V-PDB). No correction was added to the measured values, and the analytical uncertainty is estimated to be lower than 0.2‰.

Facies analysis of the Moulendé Quarry

The Moulendé quarry has been examined over an area of ~ 5 ha, for a total of nine studied outcrops (Figure 3a). Among them, the F8 outcrop (Figure 3a-c) was logged in detail because of its well-exposed strata and sedimentary structures. It is the only outcrop where the FB2a-FB2b transition is observed with accompanying black shale deposits. Lithological composition, vertical facies relationships and sedimentary features allow recognition of two

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

depositional facies from the MRS-bearing strata.

Facies 1

Thick-bedded, coarse-grained sandstones represent the main facies of the Moulendé quarry. These deposits are commonly referred to the Poubara sandstones (FB2a subunit; Figures 2, 3a-c). This facies is 15 m thick in the quarry, as well as in the closest drill core (GR5; Azziley Azzibrouck, 1986), but could be up to 100 m thick elsewhere in the basin (Weber, 1969). Interbedded thin-bedded, coarse-grained sandstones and laminated black shales form heterolithic beds. The uppermost massive sandstone bed is laterally heterogeneous in thickness and its bedding surface is undulatory. At the top of several massive sandstone beds, sedimentary features, including load structures, sandstone clasts and water escape structures, such as dishes, are present. These beds occasionally contain dm-scale hummocky cross stratification (HCS), restricted to the upper part of the succession (Figures 2, 3d; Reynaud *et al.*, 2017).

Facies 2

Numerous thin-bedded, parallel-laminated black shales with interbeds of siltstones lie conformably upon Facies 1. They form stacks of variable thickness in the upper part of the studied succession and are about 5 m thick. Connected to the underlying facies and approximately 0.4 m in height, a sandstone dyke cross-cuts these multilayered black shales (Figures 2, 3e). There is no evidence of fracturing along the dyke, indicating that the sediments were not consolidated prior to injection. The main component of facies 2 is characterized by rhythmic successions of mm-scale light and dark laminae of siltstones and shales. Small-scale cross-laminae are found within mm-thick interbedded siltstones. Load-shaped and flame-shaped structures forming convolute bedding (Figures 2, 3f) are observed in

227 interbedded medium-to-coarse-grained siltstone beds ranging from 2 to 5 cm in thickness.
228 Interference ripples are also observed, indicating paleowaves coming from two directions
229 (Figures 2, 3g). Lastly, a dm-scale bed containing dark-coloured convex laminae associated
230 with cm-scale current ripples comprise the thickest interbedded coarse-grained siltstones
231 (Figures 2, 3h).

232

233 Interpretations

234 The variation in thickness of the uppermost massive sandstone level (at ~15 m in the
235 stratigraphy) may be related to detachment folds and troughs later infilled by overlying
236 sediments (Reynaud *et al.*, 2017). Sudden and rapid deposition of coarse-grained sediments
237 produce load and dish structures by immediate dewatering (Reineck & Singh, 1980). Rapid
238 sedimentation also results in sandstone dykes formed by elevated pore pressures leading to
239 fluidization of the unconsolidated sediment. In addition, conditions that are favourable for the
240 formation of HCS are commonly found between the fair-weather and storm wave base
241 (Immenhauser, 2009). Various interpretations have been proposed for the paleoenvironmental
242 setting of the FB2a subunit. They range from a deltaic paleoenvironment subjected to tidal
243 influence (El Albani *et al.*, 2010; Ossa Ossa, 2010), to a shoreface paleoenvironmental setting
244 (Ngombi Pemba, 2014), to a turbidite lobe setting in waters deeper than 200 m (Parize *et al.*,
245 2013). Reynaud *et al.* (2017) showed that the Poubara sandstones are composed of high-
246 density sediment gravity flow deposits, emplaced during a forced regressive system tract.
247 These authors underline that the Francevillian basin was likely isolated because of the absence
248 of wave deposits. They also noticed the presence of linguoid and interference ripples that is
249 consistent with shallow water settings.

250 Thinly laminated black shales are deposited in a relatively quiet water setting but
251 interrupted by quickly deposited supplies of millimetre-to-centimetre-scale siltstones. These

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

laminae host many sedimentary structures that indicate a strong current. Reynaud *et al.* (2017) suggest that these sediments were deposited close to the maximum regressive surface whereas Parize *et al.* (2013) argue for the deposition of turbidites on an upper slope setting.

Mat-related structures (MRS) in the FB2 Member

The MRS documented here come from a 20 m thick interval of the FB2 sequence (Figure 2). Some specimens are entirely or partially pyritized, while others display a few isolated pyrite grains within the mat layers. Following the classification scheme by Sarkar *et al.* (2008), the Francevillian MRS are divided into two major categories: (1) mat-layer structures, representing potential in situ growth of a microbial mat (Figures 4-5, Figure S1) and (2) mat-protected structures (Figure 6), possibly formed by abiotic processes but requiring the protection of a microbial mat (Davies *et al.*, 2016).

1. Mat-layer structures

This group of MRS represents the “mat-layer itself that may be intact, deformed or rafted” (Sarkar *et al.*, 2008). In this work, we interpret well-preserved, intact mat growth according to the following criteria.

1a. ‘Elephant-skin’ textures

Description

‘Elephant-skin’ textures (Runnegar & Fedonkin, 1992) encompass quasi-polygonal, geometrically distinct reticulate patterns, ranging from the millimetre to centimetre scale, and bearing bulges or ridges (Figure 4a-c). They are both identified at the top of the massive Poubara sandstones and the interbedded siltstones in the black shales (Figure 2). Individual polygons are 0.3–1.3 cm in diameter, with a maximum relief of 0.2 cm.

Microscopic analyses show reticulate patterns preserved as non-homogenous dark laminae above a light, coarser material 200 to 600 μm thick (Figure 7a, f). The overlying layer consists of abundant wavy-crinkly laminae entangling 20 to 50 μm quartz grains and heavy minerals (*e.g.*, titanium oxides, apatites and monazites); the latter suggested by the relative proportions of the elements found through SEM-EDS analysis (Figure S2), as well as transmitted and reflected light microscopy. Most of the grains in the wavy-crinkly laminae are randomly arranged within a clayey matrix, and are dominated by O, Al, Si, Mg, K and Fe (Figure S2; Table S1). Reticulate patterns are also C-rich (up to 8.52%) compared to the underlying sandstone bed, as evident from the Raman (Figure 8a-c) and organic elemental analyses (Table S2). The carbonaceous layer is punctuated by tufted microstructures, either on bulge rims or within the bulges, with similar striking morphologies to each other and where clay layers are almost vertically oriented (Figures 7f, 9a-b). Based upon the description by Noffke *et al.* (2013), the height/base index is 30/100 μm to 120/220 μm long, separated at regular distances of 150 – 230 μm from each other.

Interpretation

‘Elephant-skin’ textures are formed by a specific arrangement of tufts, pinnacles, bulges and smaller-scale bulges arranged into ridges. Tufts, often arising in ridge junctions or at the intersection of microbial bulges, are proposed as a particular type-feature of cyanobacterial mat filaments growing vertically towards a light source (phototaxis) (Gerdes *et al.*, 2000; Porada & Bouougri, 2007; Bose & Chafetz, 2009). These textures are known in modern supratidal environments (Gerdes *et al.*, 2000; Gerdes, 2007; Bose & Chafetz, 2009; Taj *et al.*, 2014), but have also been recorded and described in some ancient shallow water environments from the terminal Proterozoic (Runnegar & Fedonkin, 1992; Gehling, 1999), and perhaps even the Archean (Flannery & Walter, 2012). However, Shepard & Sumner

1
2
3 302 (2010) debated the role of phototaxis in the formation of reticulate patterns and tufted
4
5 303 structures. They investigated whether filamentous cyanobacteria were influenced by light to
6
7 304 form polygonal structures or whether they were related to their own morphotype. Thus,
8
9 305 reticulate patterns may result from an undirected motility of filamentous bacteria without the
10
11 306 influence of light.
12

13 307

15 308 1b. *Putative macro-tufted microbial mat structures*

17 309 Description

19
20 310 The putative macro-tufted microbial mats are preserved as positive epireliefs that display
21
22 311 patches of vertical bump-like structures protruding 0.05–0.2 cm on the upper surface of
23
24 312 interbedded, coarse-grained siltstones (Figures 2, 4d).
25

26 313 Microscopic features seen in longitudinal thin sections reveal that the coarse-grained
27
28 314 siltstone is draped by a 50–500 µm thin, dark, clay-rich layer with floating quartz grains
29
30 315 (Figure 7b, g-h). These grains are clearly different in size compared to the coarser grains in
31
32 316 the underlying sediment. Some clay particles are micaceous and oriented more or less parallel
33
34 317 to the bedding plane. Furthermore, it can be noted that the thickness of this overlying dark
35
36 318 layer varies along the longitudinal section. The thinner dark-coloured layer is found at the top
37
38 319 of the microtopography, whereas the thicker one overlies the troughs.
39

40
41 320 SEM imaging reveals that the dark lamina encompasses heterogeneously-sized, 5–40
42
43 321 µm quartz grains and heavy minerals (Figure 9c), while clay minerals are positioned sub-
44
45 322 vertically. In addition, EDS analyses suggest that these laminated minerals are mainly made
46
47 323 of O, Si, Al, K elements, with minor Mg, Fe and Na (Table S1), consistent with the elemental
48
49 324 composition of mica and illite minerals (*e.g.* Velde & Meunier, 2008).
50

51 325

53 326 Interpretation

1
2
3 327 A similar macro-tufted microbial mat was recorded from a 2.0 Ga paleodesert environment
4
5 328 where 0.2 cm high mound-like structures are randomly arranged on a fine-grained sandstone
6
7 329 (Simpson *et al.*, 2013). The mechanism behind the development of this feature in the
8
9 330 sedimentary rock record remains unclear. However, by changing the polarity of their
10
11 331 filamentous structure, the growth of a microbial biofilm is able to produce similar bump-like
12
13 332 structures in modern sedimentary environment (Gerdes *et al.*, 2000; Gerdes, 2007; Bose &
14
15 333 Chafetz, 2009; Taj *et al.*, 2014). This pattern may also result from the direct filling of the pre-
16
17 334 existing microtopography by mat-building organisms as they grow and expand (Simpson *et*
18
19 335 *al.*, 2013). The smooth but bumpy morphology is likely enhanced by synsedimentary loading
20
21 336 and post-depositional compaction, and so this bio-sedimentary expression could be viewed as
22
23 337 a combined microbial and compaction feature.
24
25
26 338

27 28 339 1c. *Domal buildups*

29 30 340 Description

31
32 341 Domal buildups are preserved as positive bed-top features on bedding surfaces of black shales
33
34 342 (Figure 2). They are characterized by 0.1–0.4 cm convex, domal structures projecting
35
36 343 upwards from the bedding plane. Distinctive features include a cluster of low mound-shaped
37
38 344 structures, 0.4 cm in diameter, nearly equal in all instances and without any central
39
40 345 depressions (Figure 4e-f). Alternatively, they can be isolated structures with a diameter
41
42 346 ranging from 0.7 cm to 1.5 cm that seem to reveal a cauliflower-like texture associated with
43
44 347 extensive positive-relief, wrinkled structures (Figure 4g). Circular disks belonging to the
45
46 348 oldest large colonial organisms (El Albani *et al.*, 2014) seem closely associated with these
47
48 349 specific structures. Both isolated and clustered domal structures have been shown to be
49
50 350 pyritized (Table S2). Bulk sulfur level of up 22.16% within the domal structures is associated
51
52 351 with a carbon content <0.53%. There are also flat, pyritized mat-related structures without any
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

particular distinctive features that tend to develop close to the clustered domes (Figure 4f, Figure S1).

Thin sections reveal a well-defined alternation of clay and silt layers beneath the domal structures (Figure 7c). Most of the quartz particles from the underlying sediment are laminated and/or have a length of grain-to-grain contact of mostly 20–80 μm . None of these grains were found in the topmost dark, clay layer. Furthermore, the domal structures typically show an indication of internal convex lamination between all pyrite grains (Figure 7i), which reflects the position of the domes in epirelief.

Interpretation

Domal buildups have been reported from fluvial and fluviodeltaic settings in Mid-Proterozoic (Garlick, 1988; Schieber, 1998, 1999; Sheldon, 2012; Wilmeth *et al.*, 2014) to shallow water settings in Late Cambrian times (Bottjer & Hagadorn, 2007). The latter authors proposed that they “may represent different life histories, biological affinities, and/or histories of interaction with the physical environment”. Previous studies have also invoked a distinct mat growth pattern (Wilmeth *et al.*, 2014) and inferred that domal structures are built by an upward growth of microbial mats. This might reflect competition for light (Garlick, 1988; Schieber, 1999). The $\delta^{13}\text{C}_{\text{org}}$ isotopic signatures of Mesoproterozoic domal-like structures in Copper Harbor Conglomerate (Upper Peninsula of Michigan) are attributed to photosynthesis (Sheldon, 2012).

EPS possess chemical properties that protect cells by creating sharp geochemical gradients (Decho, 2000), while the anaerobic decay of the buried mat material facilitates the precipitation of diagenetic minerals such as pyrite, depending on sulfate and iron bioavailability. If this diagenetic mineralization follows individual layers within the EPS demarcated boundaries, or if it is associated with specific features, the former presence of

microbial mats are overprinted and preserved as a mineralized signature (Schieber, 1999). Thus, sulfate-reducing microorganisms probably used the carbonaceous material from the domal buildups as their electron donor, thereby releasing hydrogen sulfide to react with available iron. This mechanism is supported by the recurrent low carbon to high sulfur contents in the materials. Analyses of the sulfur isotope contained in pyrite from the Francevillian series measured by Secondary Ion Mass Spectrometry (SIMS) further revealed negative $\delta^{34}\text{S}$ values associated with widespread microbial decay, linked to the activities of the sulfate-reducing microorganisms (El Albani *et al.*, 2014).

1d. *Discoidal microbial colonies*

Description

A discoidal microbial colony (Grazhdankin & Gerdes, 2007) is a term for a mat-layer that includes disk-shaped, internal, centimetre-scale features (Figure 5a-g). Two major distinct discoidal microbial colony-like morphologies occur on bedding planes in the black shale unit (Figure 2). The first is characterized by centimetre to the decimetre scale clusters of outward-convex, rod-shaped structures with internal pyritized concentric bands, possessing a series of ~0.4 cm wide light and dark zones (Figure 5a-b). This specific feature does not exhibit any relief. The second variety is made up of well-marked circular bodies represented by sharp outlines separating the specimens from the surrounding black shale (Figure 5c-g). The disk-shaped colonies display slight internal concentric zonations, wrinkle features or a lack of any specific morphological characteristics. A variety of the specimens have a 1.1–1.7 cm high disk-shaped relief with concentric rims restricted to the edges, which may present themselves as cauliflower-like patterns varying between 3.3 and 8.2 cm in diameter (Figure 5c-e). There are numerous distinct large euhedral pyrite crystals on these structures. Less common are disk-shaped bodies of 2.3 cm in diameter and represented by a minor relief, internal

1
2
3 402 millimetre-scale wrinkles, and devoid of sharp outlines (Figure 5f). Other observable
4
5 403 discoidal structures are relatively small (~ 0.5 cm diameter) that do not have particular
6
7 404 features (Figure 5g). However, these are the only structures that are entirely pyritized by tiny
8
9 405 pyrite crystals.
10
11 406

12
13 407 Interpretation
14

15 408 Previous studies have shown that a wide range of variability exists in the morphology and
16
17 409 organization of modern discoidal microbial colonies in natural environments (Gerdes *et al.*,
18
19 410 1993; Banerjee *et al.*, 2014; Sarkar *et al.*, 2014) as well as under laboratory conditions
20
21 411 (Matsushita *et al.*, 2004). Concentric zonations with a series of light and dark bands represent
22
23 412 a particular variety of discoidal microbial colony known as ‘fairy rings’. Modern ‘fairy ring’
24
25 413 structures are ubiquitous in salt marshes (Gerdes *et al.*, 1993; Gerdes, 2007) and in shallow
26
27 414 water environments (Banerjee *et al.*, 2014). The concentric ring-shaped structures of the
28
29 415 Francevillian series discoidal structures clearly display the same striking morphologies. All
30
31 416 modern examples indicate that filamentous cyanobacteria and diatoms, which are embedded
32
33 417 in EPS and crowned by micro-pinnacles, form these concentric ring-shaped structures. This
34
35 418 arrangement is likely not random but represents an oriented behaviour that might be triggered
36
37 419 by external environmental factors (Gerdes, 2007). So far, the generally accepted mechanism
38
39 420 of formation involves escaping gas bubbles from the substrate, which may cause concentric
40
41 421 wave propagation (Gerdes *et al.*, 1994). Subsequently, chemotactic responses initiated by
42
43 422 adapted filamentous cyanobacteria may form the concentric organizational patterns. Presently,
44
45 423 the oldest discoidal microbial colonies that have been described are from Mesoproterozoic
46
47 424 and Neoproterozoic shallow submarine environments (Grazhdankin & Gerdes, 2007;
48
49 425 Banerjee *et al.*, 2010, 2014; Sarkar *et al.*, 2014), where pyritized filamentous structures are
50
51 426 sometimes preserved (Grazhdankin & Gerdes, 2007).
52
53
54
55
56
57
58
59
60

427

428 1e. *Horizontal mat growth pattern*429 Description

430 A unique, outward-convex, spindle-shaped structure extending over 34 cm is developed on
431 the bedding plane surface of the black shale unit (Figures 2, 5h). The structure appears similar
432 to the spindle-shaped ‘fairy ring’ structures described above. However, this structure is
433 represented by tiny, 0.1 cm tall, arched ridges, rising horizontally, following a well-defined
434 geometric pattern. These sharp-crested ridges of 0.15 cm in width are regularly spaced,
435 commonly dissected and slightly undulated. It appears that the laminae convexity of the
436 whole structure is unevenly distributed, ranging from 55 to 86 degrees.

437

438 Interpretation

439 The horizontal mat growth pattern could be related to ‘fairy ring’ structures with which the
440 pattern shares some similarities (*e.g.* outward-convex, spindle-shaped, sharp-crested ridges).
441 Nevertheless, there are no alternations of light and dark zones. Processes implicated in the
442 formation of ‘fairy ring’ structures may be involved here, but topography is added. The
443 orientation of the wave propagation and subsequent millimetre-sized ripples may be induced
444 by gravity. Then, pinnacle-forming organisms would interact with the nutrient front by
445 chemotactic responses to colonize the substratum, producing the distinct morphology
446 preserved in the sedimentary record. Alternatively, localized mat failure and strain induced by
447 gravity, waves or currents could also explain the morphology as a mechanical deformation of
448 biogenically stabilized sediment. Although it has not been suggested in other studies,
449 morphological features including a point of origin, persistent “growth” in one vector, and
450 sharp boundaries could also be explained by an underwater seep that is confined by a biomat
451 and promotes mat growth in a downslope direction. In any case, the parallel arrangement of

1
2
3 452 ridges, as well as the regularity of components in the horizontal mat growth pattern, suggests
4
5 453 a biotic origin.
6
7 454

9 455 2. Mat-protected structures

11 456 This category of mat-related structures arises from structures that probably require mats for
13
14 457 their preservation, the excepting being wrinkle marks.
15
16 458

18 459 2a. *Wrinkle structures*

20 460 Description

22 461 Based on morphology, three types of wrinkle structures are categorized here: wrinkle marks,
24 462 parallel wavy wrinkle structures and cross-cutting wrinkle structures. Wrinkle marks are
26 463 represented as laterally continuous or discontinuous, slight, outward-convex positive relief on
28 464 bedding plane of the black shales (Figures 2, 4h).

31 465 Parallel wavy wrinkle structures are preserved as sinuous, continuous ridges separated
33 466 by parallel, narrow valleys extending for less than 10 cm on bedding surfaces of sandstones
35 467 and interbedded thin-to-medium-bedded siltstones (Figures 2, 6a-b). The ridges have rounded
37 468 tops and steep sides, with an individual ridge elevation of 0.1 cm and a spacing of 0.4 cm.
39 469 Organic elemental analysis of one sample from the black shale unit does not reveal significant
41 470 differences in carbon content between the parallel wavy wrinkle structures and the
43 471 surrounding sediment (Table S2). However, microscopic analysis of thin sections, cut
45 472 perpendicularly through the wrinkles, indicates that a dark-coloured layer caps the coarse, silt-
47 473 sized rippled structures. Further down, the sediment displays quasi-planar laminations (Figure
49 474 7d). For the length of the section, the topmost carbonaceous layer is similar in thickness and
51 475 does not onlap the rippled bed, but instead reproduces its topography (Figure 7j). Oriented
53 476 quartz grains trapped within the carbonaceous layer (Figure 7j) appear to have the same grain

size as those found in the rippled structure. Tiny floating grains are also embedded in the clay-sized fractions. Sheets of various clay particles entangle several heavy minerals, mainly titanium oxides (Table S1).

Cross-cutting wrinkle structures (Banerjee & Jeevankumar, 2005) are characterized by bifurcating and frequently interconnected asymmetric ridges with intervening sub-parallel troughs on the bedding surface of black shales (Figures 2, 6c). The rounded ridges are 0.1 to 0.2 cm high, 0.4 to 0.5 cm wide, and their spacings vary from 0.4 to 1 cm. A few dome-like structures of 0.5 to 0.6 cm are close to these cross-cutting wrinkle structures (Figure 6c).

Interpretation

Wrinkle structures are one of the most common sedimentary features related to microbial mats. They are found over a large paleogeographic range, including the deep-sea (Buatois & Mángano, 2003; Flood *et al.*, 2014), shallow marine (Hagadorn & Bottjer, 1997; Banerjee & Jeevankumar, 2005; Sarkar *et al.*, 2006; Porada & Bouougri, 2007; Mata & Bottjer, 2009; Banerjee *et al.*, 2014; Buatois *et al.*, 2014; Sarkar *et al.*, 2014, 2016; Yang *et al.*, 2017), fluvial and lacustrine (Chu *et al.*, 2015, 2017), and even continental environments (Simpson *et al.*, 2013). Some researchers claimed that relative microbial community growth correlates to with irregular surface of wrinkle structures (Hagadorn & Bottjer, 1997; Banerjee & Jeevankumar, 2005; Sarkar *et al.*, 2006, 2014).

Loading and dewatering processes have also been proposed for wrinkle generation (Noffke *et al.*, 2002). A recent study has proposed that abiotic mechanisms are unable to create millimetre-scale ripples (3–15 mm wavelengths) because waves with small orbital amplitudes are not strong enough to mobilize sand and silt grains (Mariotti *et al.*, 2014). In wave tank experiments, millimetre-scale ripples were perfectly reproduced by means of microbial aggregates rolling along the substrate and transporting grains. The preservation of

1
2
3 502 such structures requires burial by fine material that may be represented by the biostabilization
4
5 503 of a bacterial community. The dark-coloured layer overlapping the silt-sized rippled structures
6
7 504 (Figure 7d, j) is probably induced by EPS that acts as a trap for particles from the surrounding
8
9 505 environment (Hagadorn & Bottjer, 1997). Oriented grains are result of a growing biofilm that
10
11 506 envelops, lifts, rotates and orientates the grains (Noffke *et al.*, 1997, 2001). By combining
12
13 507 morphological description and petrographic study, it appears that wrinkle structures are
14
15 508 polygenic. Both mat-layer and mat-protected structures are therefore most likely represented
16
17 509 in this case.
18
19

20 510
21
22 511 2b. ‘*Kinneyia*’ structures
23

24 512 Description
25

26 513 ‘*Kinneyia*’ structures are characterized by clearly distinct, short, sinuous ridges and troughs of
27
28 514 0.2 cm in width and less than 0.1 cm in height on the bedding surface of the black shale unit
29
30 515 (Figures 2, 6d). These structures fit with the definition of ‘*Kinneyia*’ as proposed by (Porada
31
32 516 *et al.*, 2008), *i.e.*, they are “comparatively short, curved, frequently bifurcating, flat-topped
33
34 517 crests, 0.5–1 mm high and 1–2 mm wide, which are separated by parallel, round-bottomed
35
36 518 depressions. The crests are usually steep sided and may run parallel”.
37
38
39

40 519
41
42 520 Interpretation
43

44 521 The origins of “*Kinneyia*” structures have long been amongst the most problematic to
45
46 522 understand. From their initial description as fossil algae (Walcott, 1914) to the present day
47
48 523 where the term is now used to describe MRS, they have been interpreted in various ways
49
50 524 (Hagadorn & Bottjer, 1997; Noffke *et al.*, 2002; Porada & Bouougri, 2007; Porada *et al.*,
51
52 525 2008; Thomas *et al.*, 2013; Mariotti *et al.*, 2014). However, few hypotheses properly explain
53
54 526 how the underlying sediment is affected (Davies *et al.*, 2016). Recently, ‘*Kinneyia*’ structures
55
56
57
58
59
60

were successfully reproduced in wave tank experiments using microbial aggregates (Mariotti *et al.*, 2014), although the sharply defined morphology of the depressions preserved might suggest that gas bubbles formed beneath the biomat could also be a possible mechanism (Pfluger, 1999). It has also been demonstrated that ‘Kinneyia’ is a polygenetic texture (Davies *et al.*, 2016), but the Francevillian ‘Kinneyia’ structures do not rule out formation by biotic mediators.

2c. Linear patterns

Description

Linear patterns are characterized by 0.1–0.3 cm high, parallel, linear ridges, slightly undulated on the bedding surfaces of massive sandstones and interbedded siltstones (Figures 2, 6e, g). They are of varying size, ranging from 2 to 8 cm in length and 0.3 to 0.7 cm in width. Around these linear ridges, the surface morphology is not uniformly flat but bears micrometric spots that seem to be embedded in the matrix (Figure 6f). Furthermore, Raman spectrometry indicates this dark-coloured layer is carbon-rich compared to the underlying sediments (Figure 8d-e).

Microscopic observations reveal that the 100 to 300 μm thick, carbon-rich layer lies on lighter, sand-sized quartz grains and consists of multiple, well-defined sheets of clay minerals (Figure 7e, k). SEM imaging shows that numerous small quartz grains and heavy minerals, such as apatite, titanium oxide and zircon, are embedded in the clay matrix (Figure 9d-e). This matrix is formed by O, Si, Al, Mg, Fe and K elements, *i.e.* chlorite and illite, as expressed by EDS analyses (Table S1). Furthermore, sand-sized quartz grains are observed inside the ridges as well as the surrounding sediment but are separated by a clayish undulated layer (Figure 9e). The quartz grains inside the ridges and those from the host rock appear randomly oriented.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

552

553 Interpretation

554 According to the identification of linear patterns in modern tidal flats by (Porada & Bouougri,
555 2007) “straight to irregular ridges are developed at distance of 1–2 cm and are locally
556 interrupted or reduced to faint lines on the flat mat surface”. Towards the water line, oriented
557 bacterial filaments dominate these structures. In experimental studies, Shepard & Sumner
558 (2010) observed linear ridges of up to 15 cm long made of cyanobacterial filaments running
559 parallel to the incident light. Moreover, it has been proposed that a faint rippled surface may
560 imply a linear pattern because of the preferential microbial growth following slight ridges
561 (Bouougri & Porada, 2007; Gerdes, 2007; Porada & Bouougri, 2007).

562 The presence of carbonaceous material within the dark-coloured layer is congruent
563 with a biotic origin. Microtextures, such as tiny quartz particles and heavy minerals embedded
564 in the matrix, are caused by the trapping and binding processes related to microbial activity
565 (Gerdes, 2007; Noffke, 2010). Individual grains bound in the organic matrix can be compared
566 to those observed with laboratory-grown cyanobacteria. These are described as oriented
567 grains that are pushed upwards during their growths (Noffke *et al.*, 2001; Noffke, 2010).

568 On a macroscopic level, the Francevillian MRS strongly resemble the linear structures
569 described above, commonly equivalent in shape but slightly smaller in size. However, no
570 microscopic features were described until now. Taking microtextures into account, it seems
571 unlikely that bacterial orientation is capable of mobilizing large amount of sediments to form
572 ridges. Microbial shrinkage caused by a period of subaerial exposition, is also a possibility
573 (e.g. Chu *et al.*, 2017; Kovalchuk *et al.*, 2017), although the surface morphologies and
574 microtextures are not consistent with this process. Also, dilational strain on a mildly dipping
575 sediment surface could explain these features, but their orientations are irregular and their
576 distributions are sporadic. Liquefaction underneath the microbial mat, due to a rise in pressure

(e.g. Porada *et al.*, 2007), could have been the cause of these particular ridges but neither sediment rising nor upward-facing microbial laminae beneath the ridges have been observed. Finally, linear patterns may result from two growth periods interrupted by small and rippled sedimentary structures (Figure 9e). This condition might be the best explanation for microbial laminae throughout the linear ridges, and the second mat growth period may have followed the ridge orientation as previously thought (Bouougri & Porada, 2007; Gerdes, 2007; Porada & Bouougri, 2007). Indirectly, linear patterns are considered to be mat-protected structures.

2d. Nodule-like structures

Description

Nodule-like structures are characterized by an elongated shape approximately 10 cm long and an irregular surface topography on the bedding surfaces of black shales (Figures 2, 6h). The surfaces of nodule-like structures comprise several millimetre-scale, tiny, crinkled ridges that are randomly distributed. Composed of silt-sized grains and capped by a dark-coloured wrinkle layer, the nodule-like structure was observed in a polished slab.

Interpretation

Similar nodular to biscuit-like surface structures have been observed in laboratory-cultured bacteria and in modern supratidal settings (Gerdes, 2007). This atypical morphology is thought to be produced by a relative abundance of coccoid cyanobacteria. However, the Francevillian nodular structures possibly reflect mat-protected structures instead of mat growth structures since the internal part of the nodules is not composed of organic matter but rather of quartz particles.

3. Isotopic analyses

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The $\delta^{13}\text{C}$ values (V-PDB) of the organic fraction, measured on different mat morphologies in both sandstone and black shale facies range from -30.67‰ to -41.26‰ (Table 1). These values are similar to previous determinations on the bulk $\delta^{13}\text{C}$ of the organic matter within the FB2 sequence, varying between *ca.* -35‰ to -30‰ (Gauthier-Lafaye & Weber, 2003; Canfield *et al.*, 2013).

Discussion

Biogenicity

The biogenicity criteria reviewed in Noffke (2009) and Wacey (2009) establishes the investigated mat-like structures to be of a microbial mat origin, having formed on the shallow marginal self-environment of the 2.1 Ga Francevillian basin (Reynaud *et al.*, 2017). The sedimentary facies on which the structures are identified must not have undergone metamorphism beyond greenschist grade (Noffke, 2009). This is in agreement with the absence of metamorphic overprint in the Francevillian facies (maximum temperature 100 °C; Gauthier-Lafaye & Weber, 1989; Ngombi-Pemba *et al.*, 2014). In terms of shape and size, the Francevillian MRS are a perfect match to those described in the Precambrian and the Phanerozoic (Figures S3-S4) in accordance with the size distribution between modern and ancient MRS, which should be comparable (Noffke, 2009). Typically, the taphonomic preservation window of MRS in the sedimentary record is valid, but not restricted by ecological setting (Davies *et al.*, 2016). Noffke (2009) suggested that most MRS are related to photoautotrophic mats formed in well-lit shallow marine environments, preferentially on fine sand deposits, even though recent studies argued that such features can also arise on deep marine sediments that do not receive light (Davies *et al.*, 2016). Regardless, our samples originated from a shallow marine setting associated with rapid sand flow deposits within a well-lit portion of the 2.1 Ga Francevillian continental shelf (Reynaud *et al.*, 2017).

Microtextural composition of the analysed fabric further support the biological trapping, binding (or ‘flypaper’ effect) and orientation of grains induced by the growth and development of microbial mats and their hydrologically-controlled interaction with sediments. The Francevillian MRS express this biological process by containing clay particles, floating grains of silt-sized quartz and concentrated heavy minerals, as well as by wavy-crinkly laminae (Figures 7f-k, 9, Figure S2; Table S1). These form well-defined organo-sedimentary structures caused by microbial baffling and trapping. The randomly oriented clay minerals in mat laminae suggest that they were trapped on the mat surface whereas laminated clay particles suggest a probable alignment by compaction (Schieber, 1998). Oriented grains reflect a particular microtexture (Figures 6f, 7j), while grain size matches that of the underlying substratum. It is commonly thought that these particles were dragged upwards by cyanobacterial mat growth (Noffke *et al.*, 1997, 2001).

In addition, geochemical evidence suggests bacterially induced biological processes characterized by carbonaceous material enriched in light carbon (Figure 8; Table 1) and pyritized structures (Figures 4e-g, 5a-e, g, 7c, i) that are depleted in heavy sulfur. The latter points to diagenetic mat destruction through burial decay (Noffke, 2009; Noffke *et al.*, 2013) and anaerobic respiration of that organic carbon by sulfate-reducing microorganisms (El Albani *et al.*, 2014; Hill *et al.*, 2016).

Paleoenvironmental interpretations and implications

Modern mat-related structures are mostly described in carbonate and siliciclastic environments but few have been described from shale deposits. The presence of black shales with large amounts of organic matter deposited in the photic zone may arise in restricted to isolated basins (Schwark & Frimmel, 2004). Some anoxygenic photosynthetic bacteria even prospered in these paleoecological conditions and their high productivity may be related to the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

availability of essential nutrients. For any MRS-bearing rocks, the hydraulic pattern must be moderated with a low sedimentation rate to promote the development of microbial communities on a substrate (Schieber, 1999; Gerdes *et al.*, 2000; Gerdes, 2007; Noffke, 2009, 2010). As an indicator of paleo-hydrological conditions, clay minerals and silt-sized grains within mat layers, are taken to represent currents strong enough to move thin particles but insufficient to transport sand-sized grains (Schieber, 1998; Noffke, 2009). Silt-sized sedimentary particles and heavy minerals can also be baffled and trapped by cyanobacterial filaments moving vertically upwards in order to escape being buried by the small-sized particles, as well as their need to reach optimal light conditions for growth (Noffke, 2009, 2010).

In this study, some of the mat-like structures, including the ‘elephant-skin’ morphotypes, the tufted structures, and the linear patterns, tend to be associated with silty lenses (Figure 2), suggesting that the baffling and grain trapping processes were operating in potential microbial mat-dominated environments at 2.1 Ga. Further, the comparable association of several MRS to the distribution of modern and fossil analogues have been used to improve the reliability of paleoenvironmental marine interpretations, including physical processes such as desiccation and erosion and biological activity (Bose & Chafetz, 2009; Noffke *et al.*, 2013; Banerjee *et al.*, 2014; Sarkar *et al.*, 2014). Importantly, the absence of desiccation and erosion-induced MRS in the Francevillian rocks suggest a quiet underwater environment from the time of deposition to when the sediments were lithified and buried.

The Paleoproterozoic Francevillian MRS possess a wide spectrum of morphologies that can be related to photoautotrophic microbial assemblages. Indeed, discoidal microbial colonies in the modern environments and domal buildups in ancient settings are results of cyanobacterial chemotaxis or phototaxis behaviours (Gerdes, 2007). The biological mechanisms leading to oriented grains has been reproduced with cultures of cyanobacteria

(Noffke *et al.*, 2001). Reticulate patterns and tufted microbial mats have striking morphologies with modern analogues built by filamentous cyanobacteria (Figure S3a; Gerdes *et al.*, 2000; Gerdes, 2007; Bose & Chafetz, 2009; Taj *et al.*, 2014). It is thought that their formation implies a phototactic behaviour (Gerdes *et al.*, 2000; Reyes *et al.*, 2013), whereas laboratory-cultured filamentous cyanobacteria show a dependency on oxygen concentration (Sim *et al.*, 2012). It has also been proposed that the undirected gliding motility of filamentous bacteria species may form these structures (Shepard & Sumner, 2010). Consequently, all bacteria with highly motile filaments are believed to be able to display these morphologies. Filamentous sulfur-oxidizing bacteria may also have the ability to produce ‘elephant-skin’-like and tuft-like structures (Flood *et al.*, 2014). However, research by genomic comparisons reveals that these phenotypic traits were most likely inherited by horizontal gene transfers from the cyanobacteria (Flood *et al.*, 2014). In other words, those aerobic chemolithoautotrophs may not have been in existence at 2.1 Ga. In addition, recent studies in a perennially ice-covered Antarctic lake have shown the specific assemblage of a photosynthetic microbial ecosystem (Sumner *et al.*, 2015; Jungblut *et al.*, 2016), with cyanobacteria forming tuft-related structures exclusively found in the oxic zone where irradiance is at its highest, whereas flat bacterial mats of anoxygenic photoautotrophs are restricted to the deeper euxinic zone experiencing lower irradiance. Thus, ‘elephant-skin’ and tufted structures may possibly have a link with oxygenic photoautotrophic microorganisms. Importantly, Flannery & Walter (2012) and Homann *et al.* (2015) thought that cyanobacteria are the only microorganisms capable of producing vertical structures or tufts.

Geomicrobiological implications

It is well established that carbon fixation by autotrophic organisms preferentially incorporates light ^{12}C over heavy ^{13}C isotopes in biomass (Schidlowski, 1988, 2001). In this regard, the

1
2
3 702 $\delta^{13}\text{C}_{\text{org}}$ values ranging from -30.67 to -41.26‰ (average $-32.94 \pm 1.17\text{‰}$) reported here are
4
5 703 within the expected range for autotrophic carbon fixation (Schidlowski, 1988, 2001; Berg *et*
6
7 704 *al.*, 2010). The typical $\delta^{13}\text{C}_{\text{org}}$ values generated by the widespread activity of ribulose 1,5-
8
9 705 bisphosphate carboxylase/oxygenase (RuBisCo), common in oxyphototrophic cyanobacteria
10
11 706 have average values from -20 to -30‰. These values are less negatively fractionated than
12
13 707 anoxygenic phototrophs (Quandt *et al.*, 1977; McNevin *et al.*, 2007; Berg *et al.*, 2010), while
14
15 708 more negative values $<-30\text{‰}$ are related to autotrophic carbon fixation in the reductive acetyl
16
17 709 CoA pathway (*i.e.*, methanogenesis). In the case of the latter, acetoclastic methanogenesis
18
19 710 diagenetically supplies isotopically light ^{12}C methane to anaerobic oxidation of methane
20
21 711 (AOM), the latter often comprising a consortia of sulfate reducers and methanotrophs (Conrad
22
23 712 *et al.*, 2010). Consequently, autotrophic carbon fixation via acetyl CoA pathway, combined
24
25 713 with anaerobic oxidation of fixed C_{org} , would effectively lead to deposition of residual C_{org}
26
27 714 enriched in light ^{12}C in the range found in this study. Such fixation of CO_2 coupled with
28
29 715 diagenetic recycling of phototrophically derived C_{org} would have inevitably resulted in the
30
31 716 sequential overprinting of light $\delta^{13}\text{C}$ in buried biomats. The bulk $\delta^{13}\text{C}_{\text{org}}$ signatures $<-30\text{‰}$
32
33 717 are, therefore, most parsimoniously interpreted to represent a mixed isotopic signal resulting
34
35 718 from the activities of various primary producers and heterotrophs.

36
37
38
39 719 The FB2b subunit hosts the first known multicellular organisms closely associated
40
41 720 with biomats (Figures 2, 10). Burrows, trails and resting traces of metazoans are often closely
42
43 721 associated with biomats in the past, being interpreted as sophisticated feeding behaviours (*e.g.*
44
45 722 Buatois & Mángano, 2012; Pecoits *et al.*, 2012; Buatois *et al.*, 2014; Meyer *et al.*, 2014; Chu
46
47 723 *et al.*, 2015). In modern environments, photosynthetic bacterial mats create thin O_2 -rich
48
49 724 layers, thus providing benthic O_2 oases for macroorganisms that may mine mat layers for
50
51 725 unexploited nutrients and O_2 (Gingras *et al.*, 2011). It is perhaps not a coincidence then that
52
53 726 the biomats and macroorganisms belong to the same strata. Moreover, the generation of
54
55
56
57
58
59
60

microenvironments due to the chemical properties of EPS (Decho, 2000) may permit soft-tissue mineralization, providing further protection against degradation (Sagemann *et al.*, 1999). Thus, the biomats may have stabilized the depositional surfaces and sheltered the macroorganisms, allowing them to become imprinted into the rock record. Interestingly, the large colonial organisms associated with bacterial communities are only known from the FB2b rocks. Although other black shale facies are recorded in the Francevillian basin (*e.g.*, the ~ 2.08 Ga FD black shale formation; Figure 1b), they were deposited in deeper environments beyond the euphotic zone (Canfield *et al.*, 2013).

Conclusion

Mat-related structures of the 2.1 Ga Francevillian series exhibit ten types of surface morphologies, providing a new window into the highly diversified Paleoproterozoic microbial life at that time. Mats are preserved in excellent conservation conditions from a 20 m thick interval of sandstone and black shale facies.

Microtexture analyses provide strong evidence in favour of mat-colonized sediment. Oriented grains, floating grains, heavy mineral concentrations, randomly oriented clays, pyritized structures and wavy-crinkly laminae all reflect the growth of microbial communities. Comparisons with ancient and modern analogues, as well as stable carbon isotope analyses, suggest growth within a palaeoenvironmental settings corresponding to the euphotic zone, likely less than 100m deep. Associations between Paleoproterozoic, large colonial organisms and mats may be similar to interactions of Ediacaran early metazoans and microbial carpets where O₂-producing cyanobacterial mats may explain this specific pattern. In addition, microbial mats may have played a major role in sediment biostabilization, fostering the preservation of complex macroorganisms that represent the first ecosystem comprising microbial biofilms and large colonial life forms.

1
2
3 752
4
5 753
6
7 754
8
9 755
10
11 756
12
13 757
14
15 758
16
17 759
18
19 760
20
21 761
22
23 762
24
25 763
26
27 764
28
29 765
30
31 766
32
33 767
34
35 768
36
37 769
38
39 770
40
41 771
42
43 772
44
45 773
46
47 774
48
49 775
50
51 776
52
53
54
55
56
57
58
59
60

Acknowledgements

We thank the Ministry of Mines, Oil, Energy and Hydraulic Resources; General Direction of Mines and Geology of Gabon; Ministry of Education, Research and Culture; National Center for Scientific Research of Gabon (CENAREST); Sylvia Bongo Foundation; Agence Nationale des Parcs Nationaux of Gabon; University of Masuku; COMILOG-Company; French Embassy at Libreville; and Institut Français du Gabon, French Ministry for Foreign Affairs. ECF is funded by the European Research Council (ERC) Seventh Framework Program (FP7) grant No. 336092. Funding for KK and MG were provided by the Natural Sciences and Engineering Research Council (NSERC). For information and scientific discussion, we acknowledge O. Bankole, C. Fontaine and J-Y. Reynaud. For assistance, we thank N. Guignard (Raman microprobe analysis) and J. Laduranty (CHNS analysis). In France, we are grateful to the French CNRS, FEDER, the Universities of Poitiers and Lille 1, and the Nouvelle Aquitaine Region. All co-authors agree to the publication of this manuscript and declare no conflict of interest.

777
778
779
780
781
782
783
784
785
786
787
788
789
790

References

- Azziley Azzibrouck G (1986) *Sédimentologie et géochimie du Francevillien B (Protérozoïque inférieur). Métallogénie des gisements de manganèse de Mouanda, Gabon.*
- Banerjee S, Jeevankumar S (2005) Microbially originated wrinkle structures on sandstone and their stratigraphic context: Palaeoproterozoic Koldaha Shale, central India. *Sedimentary Geology* **176**, 211–224.
- Banerjee S, Sarkar S, Eriksson PG, Hu X-F, Wang Y (2014) Palaeoenvironmental and biostratigraphic implications of microbial mat-related structures: Examples from the modern Gulf of Cambay and the Precambrian Vindhyan Basin, India. *Journal of Palaeogeography* **3**, 127–144.
- Banerjee S, Sarkar S, Eriksson PG, Samanta P (2010) Microbially related structures in siliciclastic sediment resembling Ediacaran fossils: examples from India, ancient and modern. In: *Microbial Mats: Modern and Ancient Microorganisms in Stratified Systems* (eds. Seckbach J, Oren A). Springer Netherlands, Dordrecht, pp. 109–129.
- Bankole OM, El Albani A, Meunier A, Rouxel OJ, Gauthier-Lafaye F, Bekker A (2016) Origin of red beds in the Paleoproterozoic Franceville Basin, Gabon, and implications for sandstone-hosted uranium mineralization. *American Journal of Science* **316**, 839–872.
- Bekker A, Holland HD (2012) Oxygen overshoot and recovery during the early Paleoproterozoic. *Earth and Planetary Science Letters* **317–318**, 295–304.

- 809 Bekker A, Holland HD, Wang P-L, Rumble D, Stein HJ, Hannah JL, Coetzee LL, Beukes NJ (2004)
810 Dating the rise of atmospheric oxygen. *Nature* **427**, 117–120.
- 811 Berg IA, Kockelkorn D, Ramos-Vera WH, Say RF, Zarzycki J, Hügler M, Alber BE, Fuchs G (2010)
812 Autotrophic carbon fixation in archaea. *Nature Reviews Microbiology* **8**, 447–460.
- 813 Berner RA (2004) *The Phanerozoic Carbon Cycle: CO₂ and O₂*. Oxford University Press, Oxford.
- 814 Bertrand-Sarfati J, Potin B (1994) Microfossiliferous cherty stromatolites in the 2000 Ma Franceville
815 group, Gabon. *Precambrian research* **65**, 341–356.
- 816 Bonhomme MG, Gauthier-Lafaye F, Weber F (1982) An example of Lower Proterozoic sediments:
817 the Francevillian in Gabon. *Precambrian Research* **18**, 87–102.
- 818 Bose S, Chafetz HS (2009) Topographic control on distribution of modern microbially induced
819 sedimentary structures (MISS): A case study from Texas coast. *Sedimentary Geology* **213**, 136–149.
- 820 Bottjer DJ, Hagadorn JW (2007) Mat-growth features. In: *Atlas of Microbial Mat Features Preserved*
821 *within the Siliciclastic Rock Record* (eds. Schieber J, Bose PK, Eriksson PG, Banerjee S, Sarkar S,
822 Altermann W, Catuneanu O). Elsevier, Amsterdam, pp. 53–71.
- 823 Bouougri EH, Porada H (2007) Mat-related features from the terminal Ediacaran Nudaus Formation,
824 Nama Group, Namibia. In: *Atlas of Microbial Mat Features Preserved within the Siliciclastic Rock*
825 *Record* (eds. Schieber J, Bose PK, Eriksson PG, Banerjee S, Sarkar S, Altermann W, Catuneanu O).
826 Elsevier, Amsterdam, pp. 214–222.
- 827 Bouton P, Thiéblemont D, Gouin J, Cocherie A, Guerrot C, Tegye M, Pr  at A, Simo Ndounze S,
828 Kassadou AB, Boulougui B, Ekhogha H, Moussavou M (2009) Notice explicative de la Carte
829 g  ologique de la R  publique du Gabon    1/200 000, feuille Franceville-Boumango, Libreville, 79p.
- 830 Bros R, Stille P, Gauthier-Lafaye F, Weber F, Clauer N (1992) Sm-Nd isotopic dating of Proterozoic
831 clay material: an example from the Francevillian sedimentary series, Gabon. *Earth and Planetary*
832 *Science Letters* **113**, 207–218.
- 833 Buatois LA, M  ngano MG (2003) Early colonization of the deep sea: Ichnologic evidence of deep-
834 marine benthic ecology from the Early Cambrian of northwest Argentina. *Palaos* **18**, 572–581.
- 835 Buatois LA, M  ngano MG (2012) The trace-fossil record of organism-matground interactions in space
836 and time. In: *Microbial Mats in Siliclastic Depositional Systems Through Time* (eds. Noffke N,
837 Chafetz H). SEPM (Society for Sedimentary Geology) Special Publication, Tulsa, Oklahoma, pp. 15–
838 28.
- 839 Buatois LA, Narbonne GM, M  ngano MG, Carmona NB, Myrow P (2014) Ediacaran matground
840 ecology persisted into the earliest Cambrian. *Nature Communications* **5**.
- 841 Canfield DE, Ngombi-Pemba L, Hammarlund EU, Bengtson S, Chaussidon M, Gauthier-Lafaye F,
842 Meunier A, Riboulleau A, Rollion-Bard C, Rouxel O, Asael D, Pierson-Wickmann A-C, El Albani A
843 (2013) Oxygen dynamics in the aftermath of the Great Oxidation of Earth’s atmosphere. *Proceedings*
844 *of the National Academy of Sciences* **110**, 16736–16741.
- 845 Chi Fru E, Arvest  l E, Callac N, El Albani A, Kilias S, Argyraki A, Jakobsson M (2015) Arsenic
846 stress after the Proterozoic glaciations. *Scientific Reports* **5**.
- 847 Chu D, Tong J, Bottjer DJ, Song H, Song H, Benton MJ, Tian L, Guo W (2017) Microbial mats in the
848 terrestrial Lower Triassic of North China and implications for the Permian–Triassic mass extinction.
849 *Palaeogeography, Palaeoclimatology, Palaeoecology* **474**, 214–231.

- 850 Chu D, Tong J, Song H, Benton MJ, Bottjer DJ, Song H, Tian L (2015) Early Triassic wrinkle
851 structures on land: stressed environments and oases for life. *Scientific Reports* **5**.
- 852 Conrad R, Claus P, Casper P (2010) Stable isotope fractionation during the methanogenic
853 degradation of organic matter in the sediment of an acidic bog lake, Lake Grosse Fuchskuhle.
854 *Limnology and Oceanography* **55**, 1932–1942.
- 855 Davies NS, Liu AG, Gibling MR, Miller RF (2016) Resolving MISS conceptions and misconceptions:
856 A geological approach to sedimentary surface textures generated by microbial and abiotic processes.
857 *Earth-Science Reviews* **154**, 210–246.
- 858 Decho AW (2000) Microbial biofilms in intertidal systems: an overview. *Continental Shelf Research*
859 **20**, 1257–1273.
- 860 Dubois M, Lopez M, Orberger B, Rodriguez C, Boussafir M, Dreux G, Rodrigues S, Pambo F (2015)
861 The Mn-Carbonate Rich Black Shales of the Bangombe Plateau, Francevillian Basin, Gabon. In:
862 *Mineral Ressources in a Sustainable World* (eds. André-Mayer AS, Cathelineau M, Muchez P, Pirard
863 E, Sindern S). Presented at the Proceedings of the 13th Biennial SGA Meeting, Nancy, France, pp.
864 1905–1908.
- 865 El Albani A, Bengtson S, Canfield DE, Riboulleau A, Rollion Bard C, Macchiarelli R, Ngombi Pemba
866 L, Hammarlund E, Meunier A, Moubiya Mouele I, Benzerara K, Bernard S, Boulvais P, Chaussidon
867 M, Cesari C, Fontaine C, Chi-Fru E, Garcia Ruiz JM, Gauthier-Lafaye F, Mazurier A, Pierson-
868 Wickmann AC, Rouxel O, Trentesaux A, Vecoli M, Versteegh GJM, White L, Whitehouse M, Bekker
869 A (2014) The 2.1 Ga Old Francevillian Biota: Biogenicity, Taphonomy and Biodiversity. *PLoS ONE*
870 **9**, e99438.
- 871 El Albani AE, Bengtson S, Canfield DE, Bekker A, Macchiarelli R, Mazurier A, Hammarlund EU,
872 Boulvais P, Dupuy J-J, Fontaine C, Fürsich FT, Gauthier-Lafaye F, Janvier P, Javaux E, Ossa FO,
873 Pierson-Wickmann A-C, Riboulleau A, Sardini P, Vachard D, Whitehouse M, Meunier A (2010)
874 Large colonial organisms with coordinated growth in oxygenated environments 2.1 Gyr ago. *Nature*
875 **466**, 100–104.
- 876 Eriksson PG, Sarkar S, Samanta P, Banerjee S, Porada H, Catuneanu O (2010) Paleoenvironmental
877 Context of Microbial Mat-Related Structures in Siliciclastic Rocks. In: *Microbial Mats: Modern and*
878 *Ancient Microorganisms in Stratified Systems* (eds. Seckbach J, Oren A). Springer Netherlands,
879 Dordrecht, pp. 71–108.
- 880 Farquhar J, Zerkle AL, Bekker A (2011) Geological constraints on the origin of oxygenic
881 photosynthesis. *Photosynthesis Research* **107**, 11–36.
- 882 Flannery DT, Walter MR (2012) Archean tufted microbial mats and the Great Oxidation Event: new
883 insights into an ancient problem. *Australian Journal of Earth Sciences* **59**, 1–11.
- 884 Flood BE, Bailey JV, Biddle JF (2014) Horizontal gene transfer and the rock record: comparative
885 genomics of phylogenetically distant bacteria that induce wrinkle structure formation in modern
886 sediments. *Geobiology* **12**, 119–132.
- 887 Gancarz AJ (1978) U-Pb age (2.05×10^9 years) of the Oklo uranium deposit. Presented at the The
888 Natural Fission Reactors: Annual International Atomic Energy Agency Conference, Vienna, Austria,
889 pp. 513–520.
- 890 Garlick WG (1988) Algal Mats, Load Structures, and Synsedimentary Sulfides in Revett Quartzites of
891 Montana and Idaho. *Economic Geology* **83**, 1259–1278.
- 892 Gauthier-Lafaye F (2006) Time constraint for the occurrence of uranium deposits and natural nuclear

- 893 fission reactors in the Paleoproterozoic Franceville Basin (Gabon). *Geological Society of America*
894 *Memoirs* **198**, 157–167.
- 895 Gauthier-Lafaye F, Weber F (1989) The Francevillian (Lower Proterozoic) Uranium Ore Deposits of
896 Gabon. *Economic Geology* **84**, 2267–2285.
- 897 Gauthier-Lafaye F, Weber F (2003) Natural nuclear fission reactors: time constraints for occurrence,
898 and their relation to uranium and manganese deposits and to the evolution of the atmosphere.
899 *Precambrian Research* **120**, 81–100.
- 900 Gehling JG (1999) Microbial mats in terminal Proterozoic siliciclastics; Ediacaran death masks.
901 *Palaios* **14**, 40–57.
- 902 Gerdes G (2007) Structures left by modern microbial mats in their host sediments. In: *Atlas of*
903 *Microbial Mat Features Preserved within the Siliciclastic Rock Record* (eds. Schieber J, Bose PK,
904 Eriksson PG, Banerjee S, Sarkar S, Altermann W, Catuneanu O). Elsevier, Amsterdam, pp. 5–38.
- 905 Gerdes G, Claes M, Dunajtschik-Piewak K, Riege H, Krumbein WE, Reineck H-E (1993)
906 Contribution of microbial mats to sedimentary surface structures. *Facies* **29**, 61.
- 907 Gerdes G, Klenke T, Noffke N (2000) Microbial signatures in peritidal siliciclastic sediments: a
908 catalogue. *Sedimentology* **47**, 279–308.
- 909 Gerdes G, Krumbein WE, Reineck H-E (1994) Microbial mats as architects of sedimentary surface
910 structures. In: *Biostabilization of Sediments* (eds. Krumbein WE, Paterson DM, Stal LJ). Bibliotheks
911 und Informationssystem der Carl von Ossietzky Universität Oldenburg (BIS)-Verlag, Oldenburg, pp.
912 165–182.
- 913 Gingras M, Hagadorn JW, Seilacher A, Lalonde SV, Pecoits E, Petrash D, Konhauser KO (2011)
914 Possible evolution of mobile animals in association with microbial mats. *Nature Geoscience* **4**, 372–
915 375.
- 916 Grazhdankin D, Gerdes G (2007) Ediacaran microbial colonies: Ediacaran microbial colonies. *Lethaia*
917 **40**, 201–210.
- 918 Hagadorn JW, Bottjer DJ (1997) Wrinkle structures: Microbially mediated sedimentary structures
919 common in subtidal siliciclastic settings at the Proterozoic-Phanerozoic transition. *Geology* **25**, 1047.
- 920 Hill C, Corcoran PL, Aranha R, Longstaffe FJ (2016) Microbially induced sedimentary structures in
921 the Paleoproterozoic, upper Huronian Supergroup, Canada. *Precambrian Research* **281**, 155–165.
- 922 Holland HD (2002) Volcanic gases, black smokers, and the Great Oxidation Event. *Geochimica et*
923 *Cosmochimica Acta* **66**, 3811–3826.
- 924 Homann M, Heubeck C, Airo A, Tice MM (2015) Morphological adaptations of 3.22 Ga-old tufted
925 microbial mats to Archean coastal habitats (Moodies Group, Barberton Greenstone Belt, South
926 Africa). *Precambrian Research* **266**, 47–64.
- 927 Horie K, Hidaka H, Gauthier-Lafaye F (2005) U-Pb geochronology and geochemistry of zircon from
928 the Franceville series at Bidoudouma, Gabon. Presented at the The 15th Annual Goldschmidt
929 Conference, Moscow, United States.
- 930 Immenhauser A (2009) Estimating palaeo-water depth from the physical rock record. *Earth-Science*
931 *Reviews* **96**, 107–139.
- 932 Jungblut AD, Hawes I, Mackey TJ, Krusor M, Doran PT, Sumner DY, Eisen JA, Hillman C, Goroncy
933 AK (2016) Microbial Mat Communities along an Oxygen Gradient in a Perennially Ice-Covered

- 934 Antarctic Lake. *Applied and Environmental Microbiology* **82**, 620–630.
- 935 Karhu JA, Holland HD (1996) Carbon isotopes and the rise of atmospheric oxygen. *Geology* **24**, 867–
936 870.
- 937 Konhauser KO, Lalonde SV, Planavsky NJ, Pecoits E, Lyons TW, Mojzsis SJ, Rouxel OJ, Barley ME,
938 Rosière C, Fralick PW, Kump LR, Bekker A (2011) Aerobic bacterial pyrite oxidation and acid rock
939 drainage during the Great Oxidation Event. *Nature* **478**, 369–373.
- 940 Kovalchuk O, Owttrim GW, Konhauser KO, Gingras MK (2017) Desiccation cracks in siliciclastic
941 deposits: Microbial mat-related compared to abiotic sedimentary origin. *Sedimentary Geology* **347**,
942 67–78.
- 943 Lekele Baghekema SG, Lepot K, Riboulleau A, Fadel A, Trentesaux A, El Albani A (2017) Nanoscale
944 analysis of preservation of ca. 2.1 Ga old Francevillian microfossils, Gabon. *Precambrian Research*
945 **301**, 1–18.
- 946 Lyons TW, Reinhard CT, Planavsky NJ (2014) The rise of oxygen in Earth's early ocean and
947 atmosphere. *Nature* **506**, 307–315.
- 948 Mariotti G, Pruss SB, Perron JT, Bosak T (2014) Microbial shaping of sedimentary wrinkle structures.
949 *Nature Geoscience* **7**, 736–740.
- 950 Mata SA, Bottjer DJ (2009) The paleoenvironmental distribution of Phanerozoic wrinkle structures.
951 *Earth-Science Reviews* **96**, 181–195.
- 952 Matsushita M, Hiramatsu F, Kobayashi N, Ozawa T, Yamazaki Y, Matsuyama T (2004) Colony
953 formation in bacteria: experiments and modeling. *Biofilms* **1**, 305–317.
- 954 McNevin DB, Badger MR, Whitney SM, Caemmerer GG von, Tcherkez GG, Farquhar GD (2007)
955 Differences in carbon isotope discrimination of three variants of D-ribulose-1,5-bisphosphate
956 carboxylase/oxygenase reflect differences in their catalytic mechanisms. *Journal of Biological*
957 *Chemistry* **282**, 36068–36076.
- 958 Meyer M, Xiao S, Gill BC, Schiffbauer JD, Chen Z, Zhou C, Yuan X (2014) Interactions between
959 Ediacaran animals and microbial mats: Insights from Lamonte trevallis, a new trace fossil from the
960 Dengying Formation of South China. *Palaeogeography, Palaeoclimatology, Palaeoecology* **396**, 62–
961 74.
- 962 Mouélé IM, Dudoignon P, El Albani A, Meunier A, Boulvais P, Gauthier-Lafaye F, Paquette J-L,
963 Martin H, Cuney M (2014) 2.9–1.9 Ga paleoalterations of Archean granitic basement of the
964 Franceville basin (Gabon). *Journal of African Earth Sciences* **97**, 244–260.
- 965 Ngombi Pemba L (2014) *Géochimie et minéralogie des formations argileuses (2.2 – 2.0 Ga) du bassin*
966 *de Franceville au Gabon : fluctuations de l'oxygène atmosphérique, chimie des océans et diagenèse*
967 *au Paléoprotérozoïque*.
- 968 Ngombi-Pemba L, Albani AE, Meunier A, Grauby O, Gauthier-Lafaye F (2014) From detrital heritage
969 to diagenetic transformations, the message of clay minerals contained within shales of the
970 Palaeoproterozoic Francevillian basin (Gabon). *Precambrian Research* **255**, 63–76.
- 971 Noffke N (2009) The criteria for the biogenicity of microbially induced sedimentary structures
972 (MISS) in Archean and younger, sandy deposits. *Earth-Science Reviews* **96**, 173–180.
- 973 Noffke N (2010) *Geobiology*. Springer Berlin Heidelberg, Berlin, Heidelberg.
- 974 Noffke N, Christian D, Wacey D, Hazen RM (2013) Microbially Induced Sedimentary Structures

- 975 Recording an Ancient Ecosystem in the *ca.* 3.48 Billion-Year-Old Dresser Formation, Pilbara,
976 Western Australia. *Astrobiology* **13**, 1103–1124.
- 977 Noffke N, Gerdes G, Klenke T, Krumbein WE (1997) A microscopic sedimentary succession of
978 graded sand and microbial mats in modern siliciclastic tidal flats. *Sedimentary Geology* **110**, 1–6.
- 979 Noffke N, Gerdes G, Klenke T, Krumbein WE (2001) Microbially Induced Sedimentary Structures—A
980 New Category within the Classification of Primary Sedimentary Structures: PERSPECTIVES. *Journal*
981 *of Sedimentary Research* **71**, 649–656.
- 982 Noffke N, Knoll AH, Grotzinger JP (2002) Sedimentary controls on the formation and preservation of
983 microbial mats in siliciclastic deposits: a case study from the Upper Neoproterozoic Nama Group,
984 Namibia. *Palaios* **17**, 533–544.
- 985 Ossa Ossa F-G (2010) *Etude multi-approches du bassin sédimentaire paléoprotérozoïque (2. 1-2. 4*
986 *Ga) de Franceville au Gabon: les environnements sédimentaires et l'impact des paléocirculations de*
987 *fluides.*
- 988 Parize O, Feybesse J-L, Guillocheau F, Mulder T (2013) Were the 2.1-Gyr fossil colonial organisms
989 discovered in the Francevillian basin (Palaeoproterozoic, Gabon) buried by turbidites? *Comptes*
990 *Rendus Geoscience* **345**, 101–110.
- 991 Partin CA, Lalonde SV, Planavsky NJ, Bekker A, Rouxel OJ, Lyons TW, Konhauser KO (2013a)
992 Uranium in iron formations and the rise of atmospheric oxygen. *Chemical Geology* **362**, 82–90.
- 993 Partin CA, Bekker A, Planavsky NJ, Scott CT, Gill BC, Li C, Podkovyrov V, Maslov A, Konhauser
994 KO, Lalonde SV, Love GD, Poulton SW, Lyons TW (2013b) Large-scale fluctuations in Precambrian
995 atmospheric and oceanic oxygen levels from the record of U in shales. *Earth and Planetary Science*
996 *Letters* **369–370**, 284–293.
- 997 Pecoits E, Konhauser KO, Aubet NR, Heaman LM, Veroslavsky G, Stern RA, Gingras MK (2012)
998 Bilaterian Burrows and Grazing Behavior at >585 Million Years Ago. *Science* **336**, 1693–1696.
- 999 Pfluger F (1999) Matground Structures and Redox Facies. *PALAIOS* **14**, 25.
- 1000 Porada H, Bouougri EH (2007) Wrinkle structures—a critical review. In: *Atlas of Microbial Mat*
1001 *Features Preserved within the Siliciclastic Rock Record* (eds. Schieber J, Bose PK, Eriksson PG,
1002 Banerjee S, Sarkar S, Altermann W, Catuneanu O). Elsevier, Amsterdam, pp. 135–144.
- 1003 Porada H, Bouougri EH, Ghergut J (2007) Hydraulic conditions and mat-related structures in tidal
1004 flats and coastal sabkhas. In: *Atlas of Microbial Mat Features Preserved within the Siliciclastic Rock*
1005 *Record* (eds. Schieber J, Bose PK, Eriksson PG, Banerjee S, Sarkar S, Altermann W, Catuneanu O).
1006 Elsevier, Amsterdam, pp. 258–265.
- 1007 Porada H, Ghergut J, Bouougri EH (2008) Kinneyia-type wrinkle structures - Critical review and
1008 model of formation. *PALAIOS* **23**, 65–77.
- 1009 Quandt L, Gottschalk G, Ziegler H, Stichler W (1977) Isotope discrimination by photosynthetic
1010 bacteria. *FEMS Microbiology Letters* **1**, 125–128.
- 1011 Reineck H-E, Singh IB (1980) *Depositional Sedimentary Environments*. Springer Berlin Heidelberg,
1012 Berlin, Heidelberg.
- 1013 Reyes K, Gonzalez NI, Stewart J, Ospino F, Nguyen D, Cho DT, Ghahremani N, Spear JR, Johnson
1014 HA (2013) Surface Orientation Affects the Direction of Cone Growth by *Leptolyngbya* sp. Strain C1,
1015 a Likely Architect of Coniform Structures Octopus Spring (Yellowstone National Park). *Applied and*
1016 *Environmental Microbiology* **79**, 1302–1308.

- Reynaud J-Y, Trentesaux A, El Albani A, Aubineau J, Ngombi-Pemba L, Guiyeligou G, Bouton P, Gauthier-Lafaye F, Weber F (2017) Depositional setting of the 2.1 Ga Francevillian macrobiota (Gabon): Rapid mud settling in a shallow basin swept by high-density sand flows. *Sedimentology*.
- Runnegar BN, Fedonkin MA (1992) Proterozoic Metazoan Body Fossils. In: *The Proterozoic Biosphere: A Multidisciplinary Study* (eds. Schopf JW, Klein C). Cambridge University Press, Cambridge, pp. 369–388.
- Sagemann J, Bale SJ, Briggs DE, Parkes RJ (1999) Controls on the formation of authigenic minerals in association with decaying organic matter: an experimental approach. *Geochimica et Cosmochimica Acta* **63**, 1083–1095.
- Sarkar S, Banerjee S, Samanta P, Chakraborty N, Chakraborty PP, Mukhopadhyay S, Singh AK (2014) Microbial mat records in siliciclastic rocks: Examples from Four Indian Proterozoic basins and their modern equivalents in Gulf of Cambay. *Journal of Asian Earth Sciences* **91**, 362–377.
- Sarkar S, Banerjee S, Samanta P, Jeevankumar S (2006) Microbial mat-induced sedimentary structures in siliciclastic sediments: examples from the 1.6 Ga Chorhat Sandstone, Vindhyan Supergroup, MP, India. *Journal of Earth System Science* **115**, 49–60.
- Sarkar S, Bose P, Samanta P, Sengupta P, Eriksson P (2008) Microbial mat mediated structures in the Ediacaran Sonia Sandstone, Rajasthan, India, and their implications for Proterozoic sedimentation. *Precambrian Research* **162**, 248–263.
- Sarkar S, Choudhuri A, Mandal S, Eriksson PG (2016) Microbial mat-related structures shared by both siliciclastic and carbonate formations. *Journal of Palaeogeography* **5**, 278–291.
- Schidlowski M (1988) A 3,800-million-year isotopic record of life from carbon in sedimentary rocks. *Nature* **333**, 313–318.
- Schidlowski M (2001) Carbon isotopes as biogeochemical recorders of life over 3.8 Ga of Earth history: evolution of a concept. *Precambrian Research* **106**, 117–134.
- Schieber J (1998) Possible indicators of microbial mat deposits in shales and sandstones: examples from the Mid-Proterozoic Belt Supergroup, Montana, U.S.A. *Sedimentary Geology* **120**, 105–124.
- Schieber J (1999) Microbial mats in terrigenous clastics; the challenge of identification in the rock record. *Palaios* **14**, 3–12.
- Schwark L, Frimmel A (2004) Chemostratigraphy of the Posidonia Black Shale, SW-Germany II. Assessment of extent and persistence of photic-zone anoxia using aryl isoprenoid distributions. *Chemical Geology* **206**, 231–248.
- Sheldon ND (2012) Microbially Induced Sedimentary Structures in the Ca. 1100 Ma Terrestrial Midcontinent Rift of North America. In: *Microbial Mats in Siliclastic Depositional Systems Through Time* (eds. Noffke N, Chafetz HS). SEPM (Society for Sedimentary Geology) Special Publication, Tulsa, Oklahoma, pp. 153–162.
- Shepard RN, Sumner DY (2010) Undirected motility of filamentous cyanobacteria produces reticulate mats: Motility produces reticulate mats. *Geobiology* **8**, 179–190.
- Sim MS, Liang B, Petroff AP, Evans A, Klepac-Ceraj V, Flannery DT, Walter MR, Bosak T (2012) Oxygen-Dependent Morphogenesis of Modern Clumped Photosynthetic Mats and Implications for the Archean Stromatolite Record. *Geosciences* **2**, 235–259.
- Simpson EL, Heness E, Bumby A, Eriksson PG, Eriksson KA, Hilbert-Wolf HL, Linnevelt S, Malenda HF, Modungwa T, Okafor OJ (2013) Evidence for 2.0 Ga continental microbial mats in a

- 1059 paleodesert setting. *Precambrian Research* **237**, 36–50.
- 1060 Sumner DY, Hawes I, Mackey TJ, Jungblut AD, Doran PT (2015) Antarctic microbial mats: A
1061 modern analog for Archean lacustrine oxygen oases. *Geology* **43**, 887–890.
- 1062 Taj RJ, Aref MAM, Schreiber BC (2014) The influence of microbial mats on the formation of sand
1063 volcanoes and mounds in the Red Sea coastal plain, south Jeddah, Saudi Arabia. *Sedimentary Geology*
1064 **311**, 60–74.
- 1065 Thomas K, Herminghaus S, Porada H, Goehring L (2013) Formation of Kinneyia via shear-induced
1066 instabilities in microbial mats. *Philosophical Transactions of the Royal Society A: Mathematical,*
1067 *Physical and Engineering Sciences* **371**, 20120362–20120362.
- 1068 Velde BD, Meunier A (2008) *The Origin of Clay Minerals in Soils and Weathered Rocks*. Springer-
1069 Verlag, Berlin.
- 1070 Wacey D (2009) *Early life on earth: a practical guide*. Topics in geobiology. Springer, Dordrecht.
- 1071 Walcott CD (1914) Cambrian Geology and Palaeontology III, No. 2. Precambrian, Algonkian algal
1072 flora. *Smithsonian Miscellaneous Collections* **64**, 77–156.
- 1073 Weber F (1969) *Une série précambrienne du Gabon : le Francevillien. Sédimentologie, géochimie,*
1074 *relations avec les gîtes minéraux associés*.
- 1075 Wilmeth DT, Dornbos SQ, Isbell JL, Czaja AD (2014) Putative domal microbial structures in fluvial
1076 siliciclastic facies of the Mesoproterozoic (1.09 Ga) Copper Harbor Conglomerate, Upper Peninsula of
1077 Michigan, USA. *Geobiology* **12**, 99–108.
- 1078 Yang H, Chen Z-Q, Fang Y (2017) Microbially induced sedimentary structures from the 1.64 Ga
1079 Chuanlinggou Formation, Jixian, North China. *Palaeogeography, Palaeoclimatology, Palaeoecology*
1080 **474**, 7–25.
- 1081
- 1082 **Figure captions**
- 1083 **FIGURE 1 Geological map and lithostratigraphic column.** (a) Geological map of the Francevillian basin. The
1084 studied quarry is Moulendé (green star). Geological map adopted from (Bouton *et al.*, 2009). (b) Synthetic
1085 lithostratigraphy of the Francevillian series. Four sedimentary units rest unconformably on Archean rocks. The
1086 red star indicates the detailed lithostratigraphic column observed in the Moulendé quarry (Figure 2).
- 1087 **FIGURE 2 Detailed lithostratigraphic column.** Composite columnar section of the Moulendé quarry in the
1088 FB2 unit showing the vertical distribution of ten representative types of mat-related structures (MRS) and
1089 sedimentary structures (SS).
- 1090 **FIGURE 3 Plane view and outcrop pictures of sedimentary facies in the Moulendé quarry.** (a)
1091 Representation of the quarry from plane view. Red box indicates the main studied outcrop in B (F8). F =
1092 outcrops. (b) Details of the bedding geometry at the transition between massive sandstone beds and thinly
1093 laminated black shales. (c) Closer view of B. (d) Cross-section view of decimetre-scale hummocky-cross
1094 stratifications (HCS), FB2a subunit. (e) Sandstone dyke, FB2a - FB2b transition. Coin diameter: ~ 2 cm. (f)
1095 Cross-section view of convolute structures, FB2b subunit. (g) Bedding plane view of interference ripples, FB2b
1096 subunit. (h) Longitudinal view of dark-coloured convex laminae associated with cm-scale foreset beds, FB2b
1097 subunit.

FIGURE 4 Microbial mat structures in the Francevillian B Formation (FB2): Mat-layer structures. (a-c) ‘Elephant-skin’ textures. (d) Putative macro-tufted microbial mat. (e-f) Clustered domal buildups and flat pyritized microbial structure (red arrow). Macrofossil specimens (white arrows). (g) Isolated domal buildups. (h) Wrinkle marks.

FIGURE 5 Microbial mat structures in the Francevillian B Formation (FB2): Mat-layer structures. (a-b) Discoidal mats likely representing ‘fairy ring’ structures. (c-e) Disc-shaped mats that display a cauliflower-like pattern. (f) Disc-shaped mat with internal wrinkle structures. (g) Small pyritized circular bodies. (h) Horizontal mat growth pattern.

FIGURE 6 Mat-related structures in the Francevillian B Formation (FB2): Mat-protected structures. (a-b) Parallel wavy wrinkle structures. (c) Cross-cutting wrinkle structures. (d) ‘Kinneyia’ structure. (e) Linear pattern. Dashed red box indicates the position of the magnification in f. Red arrow shows the location where the spectrometry Raman was performed. The Raman spectra is visible in Figure 8d. (f) Micrometric spots interpreting as oriented grains. (g) Linear patterns with several parallel ridges. (h) Nodular-like structure.

FIGURE 7 Optical photomicrographs of mat-related structures. (a) Transmitted thin section of ‘elephant-skin’ texture. Dashed red box denoting area magnified in f. (b) Transmitted thin section of putative macro-tufted microbial mat. Dashed red boxes denoting areas magnified in g and h. (c) Transmitted thin section of an isolated domal buildup. Dashed red box denoting area magnified in i. (d) Transmitted thin section of parallel wavy wrinkle structures. Dashed red box denoting area magnified in j. (e) Transmitted thin section of a linear pattern. Dashed red boxes denoting areas magnified in k and Figure 9e. (f) Tufted microstructures and wavy-crinkly laminae. Dashed red box denoting area magnified in Figure 9a. (g-h) Thickness variation across the mat layer with floating grains embedded by clays (red arrows). Mica (white arrows). Dashed red box denoting area magnified in Figure 9c. (i) Reflected magnified thin section of an entirely pyritized dome. An internal convex lamination is indicated by dashed red lines. (j) Clay laminae do not onlap the rippled siltstone bed but rather well follow its topography. Oriented grains (arrows). (k) High amount of quartz particles (arrows as example) within dark-coloured laminae. Dashed red box denoting area magnified in Figure 9d.

FIGURE 8 Polished slab of ‘elephant-skin’ texture and Raman spectra of both ‘elephant-skin’ texture and linear pattern. (a) Polished slab in cross-section perpendicular to bedding plane. Non-homogenous dark layer preserved above a pronounced boundary. Red arrow and white arrow indicate Raman spectra in b and c, respectively. (b) Representative Raman spectra of the microbial mat within bulges. It shows the presence of two Carbon peaks (“C”) at 1336 cm^{-1} (the “D1” disordered peak) and 1603 cm^{-1} (the “G” graphite peak). (c) Typical Raman spectra of sandstone with quartz (“Q”) peaks. (d) Representative Raman spectra of mat layers of linear pattern indicated in Figure 6e. It shows the presence of three carbon peaks (“C”) at $\sim 1170\text{ cm}^{-1}$ (“D4” disordered peak), 1344 cm^{-1} (the “D1” disordered peak) and 1603 cm^{-1} (the “G” graphite peak). (e) Typical Raman spectra of host sediment of linear pattern, with quartz (Q) peaks and very small intensities of “C” peaks.

FIGURE 9 SEM imaging of mat-related structures. (a) Magnified view of box area in Figure 7f. Upward clay laminae within tufted microstructures and wavy-crinkly layers. (b). Tufted microstructures and heavy minerals constitute bulges of the ‘elephant-skin’ texture. (c) Magnified view of box area in Figure 7h. Quartz grains, heavy minerals and randomly oriented clays constitute the dark-coloured mat layer. (d) Magnified view of box

1
2
3 1136 area in Figure 7k. Detrital particles wrapped by sheet clays. (e) Magnified view of box area in Figure 7e. Clay
4 1137 minerals above and throughout the ridge (arrow). No significant clue of liquefaction nor microbial shrinkage.
5
6 1138 **FIGURE 10 Examples of fossil macroorganisms associated with microbial mats.** (a) Pyritized lobate form
7 1139 just beneath ‘fairy ring’ structures. (b) Disk with radially striated core (arrow) lies on domal buildups. (c-d) Disk
8 1140 or lobate form and flat pyritized microbial structures on the same strata are closely associated. (e) Circular disks
9 1141 (arrows) rest on wrinkle marks. (f) Disk and lobate form are close to wrinkle marks on the same level or not.

11 1142
12
13
14 1143
15
16 1144
17
18 1145
19
20 1146
21
22 1147
23
24 1148
25
26 1149
27
28 1150
29
30 1151
31
32 1152

33
34 1153 **Captions for the supplementary information**

35
36 1154 **FIGURE S1 Additional flat pyritized microbial mats.**
37
38 1155 **FIGURE S2 EDS elemental maps of bulges of reticulate patterns in cross-section perpendicular to**
39 1156 **bedding plane.** BSE and composite elemental maps. Note the wavy-crinkly laminae with a large amount of
40 1157 embedded heavy minerals.
41
42 1158 **FIGURE S3 Photographs of mat-layer structures found in literature.** (a) Analogous ‘elephant-skin’ texture
43 1159 from modern lower supratidal, Bahar Alouane, southern Tunisia. Modified after photo published in Gerdes
44 1160 (2007). (b) Fossil reticulate pattern on bedding plane of siliciclastic beds from the Archean Tumbiana
45 1161 Formation, Australia. Modified after photo published in Flannery & Walter (2012). (c) A 2.0 billion-years-old
46 1162 tufted microbial mat from Makgabeng Formation, South Africa. Modified after photo published in Simpson *et*
47 1163 *al.* (2013). (d) Analogue clustered low mound-like structures with *Protichnites* trackways on bedding surface of
48 1164 quartz arenites from the Late Cambrian, Elk Mound Group, USA. Modified after photo published in Bottjer &
49 1165 Hagadorn (2007). (e) Modern analogous ‘fairy rings’ on soft muddy sediments from Bretagne salterns, France.
50 1166 Modified after photo published in Grazhdankin & Gerdes (2007). (f) Ancient example of outward-convex,
51 1167 spindle-shaped discoidal structures with concentric rings from the Mesoproterozoic, Sonia Sandstone, India.
52 1168 Modified after photo published in Sarkar *et al.* (2014). (g) Modern discoidal microbial colony on tidal flat from

1169 the Gulf of Cambay, India. Modified after photo published in Banerjee *et al.* (2014). Lens cap diameter: 6 cm.
1170 (h) Analogue discoidal microbial colony on bedding plane of sandstones from the Precambrian Vindhyan
1171 Supergroup, India. Modified after photo published in Banerjee *et al.* (2014).

1172 **FIGURE S4 Photographs of mat-related structures found in literature.** (a) Modern example of submerged
1173 wrinkle marks from Redfish Bay, Texas. Modified after photo published in Hagadorn & Bottjer (1997). (b)
1174 Patches of wrinkle marks on bedding surface of fine-grained sandstones from the Early Cambrian, Chapel Island
1175 Formation, Canada. Modified after photo published in Buatois *et al.* (2014). (c) Parallel wavy wrinkle structures
1176 reproduced in wave tank experiments using microbial aggregates. Modified after photo published in Mariotti *et*
1177 *al.* (2014). (d) Analogous parallel wavy wrinkle structures on bedding plane of mudstones from the Early
1178 Cambrian, Northwest Argentina. Modified after photo published in Buatois & Mángano (2003). (e) Minute
1179 “Kinneyia” structures formed with microbial aggregates in wave tank experiments. Modified after photo
1180 published in Mariotti *et al.* (2014). (f) Ancient analogue “Kinneyia” structures on bedding surface of siltstones
1181 from the Cambrian, Oeland, Sweden. Modified after photo published in Porada & Bouougri (2007). (g) Linear
1182 features from modern tidal flats of Bhar Alouane, southern Tunisia. Modified after photo published in Porada &
1183 Bouougri (2007). (h) Putative linear patterns on bedding surface of fine-grained quartzites from the
1184 Neoproterozoic Katanga Supergroup, Zambia. Modified after photo published in Porada & Bouougri (2007).




Figure 1

157x130mm (300 x 300 DPI)




Figure 2

180x197mm (300 x 300 DPI)




Figure 3

127x170mm (300 x 300 DPI)




Figure 4

151x211mm (300 x 300 DPI)




Figure 5

149x201mm (300 x 300 DPI)



Figure 6

142x185mm (300 x 300 DPI)




Figure 7

152x176mm (300 x 300 DPI)




Figure 8

121x81mm (300 x 300 DPI)




Figure 9

151x132mm (300 x 300 DPI)




Figure 10

152x151mm (300 x 300 DPI)

TABLE 1 $\delta^{13}\text{C}$ values of organic matter in mat-related structures (MRS).

Subunits	Samples	$\delta^{13}\text{C}_{\text{org}}$ (‰)
FB2b Black shale	MRS_1	-34.92
	MRS_2	-34.41
	MRS_3	-32.45
	MRS_4	-41.26
FB2a Sandstone	MRS_5	-31.68
	MRS_6-1	-32.72
	MRS_6-2	-33.32
	MRS_7	-33.95
	MRS_8	-33.55
	MRS_9	-33.66
	MRS_10	-33.61
	MRS_11	-32.28
	MRS_12	-30.67
	MRS_13	-31.95
	MRS_14	-32.03

Supplementary information

FIGURE S1




FIGURE S1 Additional flat pyritized microbial mats.

FIGURE S2




FIGURE S2 EDS elemental maps of bulges of reticulate patterns in cross-section perpendicular to bedding plane. BSE and composite elemental maps. Note the wavy-crinkly laminae with a large amount of embedded heavy minerals.


FIGURE S3

FIGURE S3 Photographs of mat-layer structures found in literature. (a) Analogous ‘elephant-skin’ texture from modern lower supratidal, Bahar Alouane, southern Tunisia. Modified after photo published in Gerdes (2007). (b) Fossil reticulate pattern on bedding plane of siliciclastic beds from the Archean Tumbiana Formation, Australia. Modified after photo published in Flannery & Walter (2012). (c) A 2.0 billion-years-old tufted microbial mat from Makgabeng Formation, South Africa. Modified after photo published in Simpson *et al.* (2013).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(d) Analogue clustered low mound-like structures with *Protichnites* trackways on bedding surface of quartz arenites from the Late Cambrian, Elk Mound Group, USA. Modified after photo published in Bottjer & Hagadorn (2007). (e) Modern analogous ‘fairy rings’ on soft muddy sediments from Bretagne salterns, France. Modified after photo published in Grazhdankin & Gerdes (2007). (f) Ancient example of outward-convex, spindle-shaped discoidal structures with concentric rings from the Mesoproterozoic, Sonia Sandstone, India. Modified after photo published in Sarkar *et al.* (2014). (g) Modern discoidal microbial colony on tidal flat from the Gulf of Cambay, India. Modified after photo published in Banerjee *et al.* (2014). Lens cap diameter: 6 cm. (h) Analogue discoidal microbial colony on bedding plane of sandstones from the Precambrian Vindhyan Supergroup, India. Modified after photo published in Banerjee *et al.* (2014).


FIGURE S4

FIGURE S4 Photographs of mat-related structures found in literature. (a) Modern example of submerged wrinkle marks from Redfish Bay, Texas. Modified after photo published in Hagadorn & Bottjer (1997). (b) Patches of wrinkle marks on bedding surface of fine-grained sandstones from the Early Cambrian, Chapel Island Formation, Canada. Modified after photo published in Buatois *et al.* (2014). (c) Parallel wavy wrinkle structures reproduced in wave tank

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

experiments using microbial aggregates. Modified after photo published in Mariotti *et al.* (2014). (d) Analogous parallel wavy wrinkle structures on bedding plane of mudstones from the Early Cambrian, Northwest Argentina. Modified after photo published in Buatois & Mángano (2003). (e) Minute “Kinneyia” structures formed with microbial aggregates in wave tank experiments. Modified after photo published in Mariotti *et al.* (2014). (f) Ancient analogue “Kinneyia” structures on bedding surface of siltstones from the Cambrian, Oeland, Sweden. Modified after photo published in Porada & Bouougri (2007). (g) Linear features from modern tidal flats of Bhar Alouane, southern Tunisia. Modified after photo published in Porada & Bouougri (2007). (h) Putative linear patterns on bedding surface of fine-grained quartzites from the Neoproterozoic Katanga Supergroup, Zambia. Modified after photo published in Porada & Bouougri (2007).

For Review Only

175 **TABLE S1**

176

Samples	Si	Al ^{IV}	Al ^{VI}	Al _{Tot}	Mg	Fe	Ti	Mn	K	Na	Ca	Σ Oct	Int
EST	2.78	1.22	1.58	2.8	0.73	0.11	0	0	0.33	0.13	0	2.42	0.47
	3.13	0.87	1.69	2.56	0.15	0.12	0.02	0	0.71	0.15	0	1.98	0.86
	2.89	1.11	1.47	2.58	0.33	0.05	0	0	0.47	0.18	0	1.84	0.65
	3.13	0.87	1.91	2.78	0.11	0	0	0	0.74	0.1	0	2.02	0.84
	2.88	1.12	1.6	2.72	0.52	0.17	0	0	0.52	0.11	0	2.29	0.63
	3.11	0.89	1.94	2.83	0.09	0	0	0	0.79	0.1	0	2.03	0.89
	2.63	1.37	1.17	2.54	0.72	0	0.26	0	0.26	0.17	0	2.15	0.43
PTMM	3.54	0.46	1.79	2.25	0.06	0.04	0	0	0.65	0.06	0	1.89	0.71
	3.41	0.59	1.88	2.47	0.1	0	0	0	0.69	0.07	0	1.98	0.76
	3.34	0.66	1.76	2.41	0.15	0.04	0	0	0.71	0.09	0.03	1.95	0.87
	3.46	0.54	1.69	2.23	0.28	0.06	0	0	0.55	0.09	0	2.02	0.64
	3.61	0.39	1.8	2.19	0.14	0	0	0	0.47	0.17	0	1.94	0.64
	3.35	0.65	1.6	2.25	0.3	0.09	0.01	0.02	0.58	0.12	0.03	2.02	0.73
	3.16	0.84	1.77	2.61	0.14	0.04	0.02	0.02	0.48	0.29	0.04	1.98	0.8
	3.06	0.94	1.56	2.5	0.16	0.25	0.02	0.01	0.86	0.07	0.01	2	0.94
	2.98	1.02	1.83	2.85	0.08	0.04	0.01	0.01	0.89	0.03	0.04	1.98	0.95
	3.03	0.97	1.79	2.76	0.12	0.04	0.01	0.02	0.75	0.11	0.04	1.98	0.9
	3.1	0.9	1.83	2.72	0.1	0.04	0.02	0.02	0.39	0.35	0.03	2	0.77
PWWS	3.17	0.83	1.39	2.22	0.46	0.4	0	0	0.41	0.12	0	2.25	0.53
	3.44	0.56	1.66	2.21	0.3	0.13	0	0	0.45	0.14	0	2.09	0.59
	3.56	0.44	1.67	2.11	0.24	0.1	0	0	0.53	0.09	0	2.02	0.62
	3.38	0.62	1.27	1.89	0.53	0.4	0.03	0	0.34	0.09	0	2.23	0.43
	3.17	0.83	1.91	2.75	0.12	0	0	0	0.75	0.1	0	2.03	0.85
	3.82	0.18	1.77	1.94	0.15	0	0	0	0.5	0.08	0	1.92	0.57
	3.1	0.9	1.76	2.66	0.18	0.07	0.01	0.02	0.62	0.14	0.02	2.03	0.78
	3.04	0.96	1.5	2.46	0.17	0.23	0.07	0.02	0.79	0.09	0.02	1.99	0.89
	3.18	0.82	1.62	2.44	0.32	0.15	0.01	0.02	0.47	0.18	0.02	2.12	0.68
LP	3.07	0.93	1.39	2.32	0.48	0.3	0.02	0.03	0.55	0.08	0.01	2.22	0.65
	2.97	1.03	1.39	2.42	0.47	0.31	0.04	0.03	0.53	0.13	0.02	2.24	0.67
	3.19	0.81	1.85	2.66	2.38	1.64	0.12	0	0.1	0.18	0	5.99	0.28
	3.31	0.69	2.12	2.81	2.48	1.98	0	0	0.02	0.06	0	6.57	0.08
	3.41	0.59	2.25	2.83	2.37	1.64	0.12	0	0.15	0.07	0	6.37	0.22
	3.06	0.94	2.48	3.42	1.98	2.07	0	0	0.2	0.08	0	6.53	0.28
	2.95	1.05	1.5	2.56	0.41	0.29	0.03	0	0.65	0.09	0	2.23	0.74
	3.08	0.92	1.88	2.8	0.08	0.06	0	0	0.77	0.18	0	2.02	0.95
	3.19	0.81	1.82	2.62	0.1	0.06	0	0	0.86	0.05	0	1.98	0.91
	3.11	0.89	0.99	1.88	0.43	0.54	0.09	0.05	0.44	0.05	0.06	2.1	0.55
	3.72	0.28	1.44	1.72	0.12	0.1	0.11	0.03	0.48	0.12	0.03	1.8	0.63

177

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE S1 Representative structural formulae of clays from microbial mats studied with EDS. EST, ‘elephant-skin’ texture; PTMM, putative tufted microbial mat; WS, wrinkle structure; LP, linear pattern.

TABLE S2

Samples	Mass (mg)	%C	%S
EST_mat	10.63	8.62	0.25
EST_host rock	16.62	0.58	0.05
DB_mat	15.32	0.53	22.16
DB_host rock	13.24	4.59	0.22
PWWS_mat	10.37	1.69	0.00
PWWS_host rock	14.17	2.2	0.00
sample4_mat	16.08	1.86	0.06
sample4_host rock	17.53	1.77	0.09
sample5_mat	12.58	1.25	0.54
sample5_host rock	13.18	2.11	0.11

TABLE S2 Organic elemental analyses (carbon and sulfur) on five microbial mats and their host sediments from the FB2 Formation. EST, ‘elephant-skin’ texture; DB, domal buildup; PWWS, parallel wavy wrinkle structure.